

NON-CROP USES AROUND THE FARMSTEAD - CHEMICAL MOWING - TALL FESCUE, ETC.

General Information

USE INFORMATION

This product is a postemergence, systemic herbicide with no soil residual activity.

It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Time to Symptoms

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds effects may not be visible for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds

Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the \"ANNUAL WEEDS\", \"PERENNIAL WEEDS\" and \"WOODY BRUSH AND TREES RATE SECTIONS\" for specific weed recommendations.

Always use the higher product application rate within the specified range when weed growth is heavy or dense or when weeds are growing in an undisturbed (non-cultivated) area.

Reduced weed control may result from treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions.

Cultural Considerations

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

Rainfastness

Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage

For best results, spray coverage should be uniform and complete. Do not spray foliage to the point of runoff.

Mode of Action

The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

No Soil Activity

Weeds must be emerged at the time of application to be controlled by this product.

Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by this herbicide and will continue to grow.

Biological Degradation

Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates

The maximum application or use rates stated throughout this product's labeling are given in units of volume (fluid ounces, pints, or quarts) of this product per acre.

However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as tank mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate does not exceed the maximum allowed. See the \"INGREDIENTS\" section of the label for necessary product information.

Except as otherwise specified in a crop section of the label, the combined total of all treatments must not exceed 8 quarts of this product (6 pounds of glyphosate acid) per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product (8 pounds of glyphosate acid) per acre per year.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with the label may result in injury to persons, animals or crops, or have other unintended consequences.

WEED RESISTANCE MANAGEMENT

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on

the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using other cultural or mechanical practices.

Weed Management Recommendations

To minimize the occurrence of glyphosate-resistant biotypes observe the following weed management recommendations:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your county extension agent.

Management Recommendations

For Glyphosate Resistant Biotypes

Note: Appropriate testing is critical in order to determine if a weed is resistant to glyphosate. Contact your local county extension agent or visit the following websites www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of the label.

Since the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Tacoma Ag, LLC is not responsible for any losses that may result from the failure of this product to control glyphosate resistant weeds biotypes.

The following good agronomic practices are specified to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system. This product may be applied with the following application equipment:

Aerial-Fixed wing and helicopter.

Ground Broadcast Spray-Boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held or Backpack Spray Equipment--Backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other handheld and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment-- Shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems--Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA)--Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING THE DESIRED VOLUMES.

Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED IN THE LABEL.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling. Use the specified rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on the label. Unless otherwise specified, do not exceed 2 quarts per acre. Refer to the individual use area sections of the label for further instructions.

This product, when tank mixed with dicamba, may not be applied by air in California.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be used for aerial application in California. Tank mixtures with 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and for alfalfa and pasture renovation applications only.

Ensure uniform application - to avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may help prevent corrosion.

FOR AERIAL APPLICATION IN CALIFORNIA ONLY

Extreme care must be exercised to avoid contact of spray with foliage, green stems, or fruit of desirable crops, plants, trees, or other desirable vegetation since severe damage or destruction may result.

Noncrop sites

When applied as directed and under the conditions described in the “Weeds Controlled” section of the label booklet for this product, this herbicide will control or partially control the labeled weeds growing in the following industrial, recreational and public areas, including airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, manufacturing sites, office complexes, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, roadsides (guardrails, shoulders), schools, storage areas, utility substances, warehouse areas and other public sites.

AVOID DRIFT – DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure above the manufacturer's recommendation.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application - to avoid streaking and uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may prevent corrosion.

For Aerial Application in Fresno County, California Only

From February 15 through March 31 only. For aerial application outside of these dates, refer to the "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" section printed above.

Applicable Area

This supplement only applies to the area contained inside the following boundaries

within Fresno County, California.

North: Fresno County line

South: Fresno County line

East: State Highway 99

West: Fresno County line

Use Information

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this herbicide. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Recommendations

A written recommendation MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application.

This written recommendation MUST state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and the label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this herbicide is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides.

All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-in constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the

Fresno County Agricultural Commissioner.

Application at night - Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

Read the "WARRANTY" section in the label booklet before using this product.

Ground Broadcast Equipment

Apply the product rates specified in the label in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the specified range to ensure complete coverage. Carefully select proper nozzles to avoid generating a fine mist. For best results with ground application equipment, use flat spray nozzles. Check spray pattern for uniform distribution.

Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled on a spray-to-wet basis; do not spray to the point of run-off. Spray coverage should be uniform and complete.

Use coarse sprays only. For specific rates and timing, refer to the "ANNUAL WEEDS -- HAND-HELD OR BACKPACK EQUIPMENT" section of this product label.

For low-volume directed spray applications, spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

Selective Equipment

This product may be diluted in water and applied through shielded sprayers, hooded sprayers, wiper applicators or sponge bars to weeds growing in any non-crop site specified on the label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants), and wipers may be used over-the-top of crops only when specified in this product's labeling. Such equipment must be

capable of preventing all crop contact with the herbicide solutions and operated without leakage of spray mists or dripping onto crop.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of this product with desirable vegetation may result in unwanted plant damage or destruction. Crop injury may occur when the foliage of treated weeds comes into contact with leaves of the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected.

This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Sprayers

When applied as directed under conditions described for shielded applicators, this product will control those weeds listed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of the label. Use the following equation to convert from a broadcast rate per acre to a band rate per acre.

A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Adjust the shields on these sprayers to protect desirable vegetation.

When applying to crops grown on raised beds, ensure that the hood is capable of completely enclosing the spray pattern. If necessary, extend the front and rear flaps of the hooded applicator downward to reach the ground in deep furrows. **EXTREME CARE MUST BE TAKEN TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be configured and operated in a manner that minimizes bouncing and avoids raising the hoods up off the ground at any time. If the hood is raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operating this equipment on rough or

sloping terrain where the spray hoods might be raised up off the ground surface.

Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products for use in injection systems.

CDA Equipment

The rate of this product applied per acre by vehicle-mounted controlled droplet applicator (CDA) equipment must not be less than the amount specified in the label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply in 2 to 15 gallons of water per acre.

For control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet applicators produce a spray pattern that is not easily visible. Extreme care must be taken to avoid spray or drift onto the foliage or any other green tissue of desirable vegetation, as damage or destruction of the plant may result.

ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES DIRECTIONS THAT APPLY TO ALL CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the \"ROUNDUP READY CROPS\" section of the label or separately published Supplemental Labeling for this product for instructions on treating Roundup Ready crops.

USE INSTRUCTIONS:

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed in the label, except where specifically limited. For any crop not listed in the label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the \"ANNUAL WEEDS\", \"PERENNIAL WEEDS\", AND \"WOODY BRUSH AND TREES RATE SECTIONS\" in the label. Specific rates of this product that are emphasized in this product's labeling to control tough weeds take precedence over the rates in the \"ANNUAL WEEDS\" and \"PERENNIAL WEEDS RATE SECTIONS\". Repeat applications may be made up to a maximum of 8 quarts of this product per acre per year.

Post-directed hooded sprayers and wiper applicators capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted in the individual crop sections that follow, wipers may also be used above certain crops to control tall weeds. Refer to the \"SELECTIVE EQUIPMENT\" section of the label for essential precautions regarding crop injury. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified in this product's labeling. Refer to the \"AERIAL EQUIPMENT\" section of the label for additional information.

PRECAUTIONS, RESTRICTIONS:

Avoid contact of this herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops as severe crop injury or destruction may result. When making at-planting and preemergence applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death of emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any crops not listed on the label. See \"APPLICATION EQUIPMENT AND TECHNIQUES\" section of the label for additional

information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

For broadcast postemergence treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

ROUNDUP READY CROPS

The following instructions or those published separately on supplemental labeling for this product include all applications of this product that can be made onto the specified Roundup Ready crops during the complete cropping season. DO NOT combine these treatment instructions with those for crops in the \"ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)\" section of the label that do not contain the Roundup Ready gene, unless otherwise directed in this product's labeling.

Tacoma Ag, LLC INTENDS USE OF THIS PRODUCT FOR POSTEMERGENCE (INCROP) APPLICATION ONLY ON CROPS DESIGNATED AS CONTAINING THE ROUNDUP READY GENE. Applying this product to crops that are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain the Roundup Ready gene, since severe plant injury or destruction will result.

The Roundup Ready designation indicates that the crop contains a patented gene that provides tolerance to this product. Information on Roundup Ready crops may be obtained from your seed supplier. Roundup Ready crops must be purchased from an authorized licensed seed supplier.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground broadcast applications: Apply this product in 5 to 20 gallons of spray

solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications: All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling, in particular in the \"Aerial Equipment\" section. Apply this product in 3 to 15 gallons of water per acre. See the \"APPLICATION EQUIPMENT AND TECHNIQUES\" section of the label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment.

Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

NOTE: The following instructions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burn-down treatment of this product is specified to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed with multiple germination times, or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Rates of this product specified in the following sections, or in separate supplemental labeling for this product on the control of tough weeds, supersede the rates specified in the \"ANNUAL WEEDS RATE SECTION\" and the \"PERENNIAL WEEDS RATE SECTION\" of the label.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

Chemical Mowing

USE INSTRUCTIONS: This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

PRECAUTIONS, RESTRICTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Method

[Chemical Mowing](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)