

## **ORNAMENTAL BULBS - WEEDS CONTROLLED AT 150 LBS PER ACRE (3.5 LBS PER 1000 SQ FT)**

### General Information

### General Information

Snapshot DG specialty herbicide is a preemergence product for control of certain broadleaf weeds and annual grasses in container and landscape ornamentals, nursery stock, Christmas trees, groundcovers/perennials, non-bearing fruit and nut trees, non-bearing vineyards and non-cropland areas. Apply Snapshot DG anytime prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with rate of Snapshot DG applied, weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed.

### General Use Precautions and Restrictions

In Arizona: The state of Arizona has not approved Snapshot DG for use on plants grown for agricultural/commercial production; such as on Christmas tree plantations, container grown or field grown ornamentals, non-bearing fruit and nut trees and non-bearing vineyards.

Snapshot DG controls weeds growing from seed. Snapshot DG does not control established weeds, weeds growing from stolons, rhizomes, or root pieces. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation is required to activate Snapshot DG. Optimum weed control is obtained when Snapshot DG is activated within 3 days of application. If rainfall or irrigation does not occur within 3 days of application and tillage is possible, Snapshot DG may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1 to 2 inches of soil. Failure to activate Snapshot DG within 3 days of application may result in erratic control of annual grasses. Do not apply when wind conditions favor drift of Snapshot DG granules from the target area.

Repeat applications at 150 lb per acre and higher should not be made sooner than 60 days after a previous application of Snapshot DG. Do not apply over 600 pounds per acre total of Snapshot DG within a 12-month period.

Tolerance of Turfgrass Adjacent to Ornamental Plantings: Accidental application to turf may occur when applying Snapshot DG to ornamental plantings. Snapshot DG is not recommended for weed control in turfgrasses, but turfgrasses are generally tolerant to small amounts of this product that fall outside of the intended area of application.

Do not aerially apply Snapshot DG.

Treatment Species Not Listed on the Label for Snapshot DG

Users who wish to use Snapshot DG on plant species not recommended on the label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury for during 30 to 60 days of normal growing conditions to determine if the treatment is safe to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Snapshot DG on plant species not recommended on the label.

Application Instructions

Apply Snapshot DG using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or crop injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

#### Application Techniques for Applying Snapshot DG

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.
- The weight-to-volume conversion table below provides equivalent amounts of Snapshot DG.

#### Limitations, Restrictions, and Exceptions

##### Ornamental Bulbs

Do not use Snapshot DG in bulb production fields. For use in landscape settings only.

Snapshot DG may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Snapshot DG to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual

weeds. Snapshot DG may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Snapshot DG again in late winter or early spring to weed-free soil surfaces.

#### Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.

Do not apply to bulbs while they are flowering.

#### Method

[Spray](#)

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

#### Restricted Entry Interval

12 hours

#### Timings

[Preemergence \(Weed\)](#)