

COMMERCIAL SOD PRODUCTION - NEWLY PLANTED PERENNIAL RYEGRASS AND TALL FESCUE GROWN FOR SOD

General Information

USE PRECAUTIONS

Willowood Ethofumesate 4SC or tank mixes must be used for label listed purposes and at label specified rates only. (DO NOT OVERTREAT). Do not graze livestock on treated crops.

Do not apply this product through any type of irrigation system.

If crop is lost due to climatic or soil conditions following application of Willowood Ethofumesate 4SC or tank mixes, do not plant crops other than sugar beets or ryegrass in treated land during the same season. Do not retreat field with Willowood Ethofumesate 4SC. If fields are replanted to sugar beets, reseed into treated band.

Do not rotate with any crops other than sugar beets or ryegrass for:

- 12 months following preplant incorporated, preemergence, conventional postemergence applications, or split (low rate) applications totaling more than 12 fl. oz. (0.375 lb. ai/acre);
- 6 months following split (low rate) postemergence applications totaling 12 fl. oz. (0.375 lb. ai/acre) or less.

Thorough tillage, including moldboard plowing, should precede the planting of crops other than sugar beets or ryegrass. Do not use Willowood Ethofumesate 4SC on muck or peat soils.

Do not allow spray mixture to stand in tank overnight. Flush and drain spray equipment after each day's use. Store unused spray mixture in tightly-sealed containers and protect from frost.

The label must be in the possession of the user at the time of pesticide application.

Limitations, Restrictions, and Exceptions

COMMERCIAL SOD PRODUCTION

(For use in California, Idaho, Nevada, Oregon, and Washington only)

PRODUCT INFORMATION Willowood Ethofumesate 4SC is a selective herbicide for use in established and newly planted tall fescue and perennial ryegrass grown for sod in California, Idaho, Nevada, Oregon, and Washington. Willowood Ethofumesate 4SC may be applied preemergence or postemergence for the control of weed species listed below. Overhead irrigation or rainfall shortly after application is necessary for activation.

Do not harvest treated sod for 16 days following application.

Residual control of weeds is dependent upon soil moisture conditions; rate of herbicide used, and soil texture. The activity of Willowood Ethofumesate 4SC in the soil is reduced as soil texture becomes finer and organic matter/ thatch increases.

Spray equipment: Use a fixed-boom power sprayer properly calibrated to a constant speed and rate of delivery. Do not use smaller than a 50-mesh strainer. Avoid overlapping of spray swath. Shut off boom while starting, turning, or stopping to avoid over-application. Make applications in 10 to 50 gallons of water per acre at low pressure (20 to 50 psi).

Soil preparation: All existing vegetative matter should be thoroughly worked into the soil surface before planting. Large clods, trash, or vegetative matter left on the soil surface will reduce effectiveness of the Willowood Ethofumesate 4SC treatment.

NEWLY PLANTED PERENNIAL RYEGRASS AND TALL FESCUE GROWN FOR SOD

Apply Willowood Ethofumesate 4SC to newly planted areas when crop reaches the 2- to 3-leaf stage of growth. For best results, apply to moist soils.

Before weed emergence: Apply Willowood Ethofumesate 4SC at 2 1/4 to 3 pints per acre prior to weed emergence. Use the higher rate where rattail fescue, wild oats, and volunteer cereals are expected to be a problem.

After weed emergence: Apply Willowood Ethofumesate 4SC at earliest possible weed growth stage but no later than the 4-leaf stage. Rattail fescue, wild oats, and volunteer cereals which are more difficult to control, must be treated no later than the 2-leaf stage. Apply Willowood Ethofumesate 4SC at 2 1/4 to 3 pints per acre.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Soils

[Medium](#)

[Fine](#)

Timings

[Preemergence \(Weed\)](#)

[Postemergence \(Weed\)](#)