

TURF GROWN FOR TURF OR SEED - POSTEMERGENCE - SEED FORMING STAGES

General Information

PRODUCT INFORMATION

Willowood Pronamide 3.3SC is effective for the control of a wide range of grasses and certain broadleaf weeds. The product is a soil active herbicide with uptake by sensitive weeds occurring through the roots. Before using this herbicide for a specific crop use, study the following product use information that provides important instructions for the safe and effective application of the product.

Use Restrictions: Hand-spray applications of Willowood Pronamide 3.3SC may be made only to ornamentals and nursery stock of ornamentals.

Chemigation: Do not apply this product through any type of irrigation system except as specified on the label or in Willowood, LLC supplemental labeling.

SPRAY DRIFT MANAGEMENT (AERIAL APPLICATION)

Avoiding spray drift at the application site is the responsibility of the applicator. The potential for spray drift is determined by the interaction of many equipment-and-weather-related factors. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where certain states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory Information section.

Aerial Spray Drift Advisory Information

This section is advisory in nature and does not supersede mandatory label requirements.

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity and Temperature Inversion section of the label).

Controlling Droplet Size:

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer’s specified pressures. Use the lower spray pressures specified for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets and lower drift than other nozzle types.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Do not make applications at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase swath adjustment distance with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Avoid application below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Do not apply during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. The cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. The presence of inversion conditions can be indicated by ground fog. However, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversion conditions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: Apply this pesticide when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

RESISTANCE MANAGEMENT

Willowood Pronamide 3.3SC is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Willowood Pronamide 3.3SC will not control known Group 3 resistant biotypes of labeled weeds. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance consider:

- Where possible, rotate the use of Willowood Pronamide 3.3SC or other Group 3 herbicides with different herbicide groups that control the same weeds in a field.
- For best resistance management stewardship, avoid use more than once per season and use Willowood Pronamide 3.3SC in programs with other herbicides with different modes of action.
- Where possible, rotate the use of Willowood Pronamide 3.3SC or other Group 3 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based upon an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistance weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisers for any additional pesticide resistance management and/or integrated weed management requirements for specific crops and weed biotypes.

DOSAGE

The rate of Willowood Pronamide 3.3SC required will vary depending on the crop culture involved and weed species to be controlled. See specific crop use directions for all dosage instructions. All dosage instructions listed in the label are in terms of pints of product or pounds of active ingredient per broadcast acre. For banded application, reduce the amount of Willowood Pronamide 3.3SC used per square acre according to the given formula.

TIMING AND APPLICATION

Unless specific directions are given under the crop to be treated, apply Willowood Pronamide 3.3SC in the fall or early winter, when temperatures do not exceed 55°F, but prior to freeze-up. Best weed control results occur when Willowood Pronamide 3.3SC is applied preemergence to the weeds and when application is followed by rainfall or irrigation to move the product into the root zone of the germinating weeds.

Mix Willowood Pronamide 3.3SC thoroughly in clean water at the required concentration and apply uniformly as a spray.

For ground application, use a conventional low-pressure herbicide sprayer equipped with flat fan nozzles spaced and calibrated to uniformly deliver 20 to 50 gallons of spray per acre. For aerial applications, apply in a coarse droplet spray at 5 to 10 gallons per acre. Accurately calibrate spray equipment prior to each use.

COMPATIBILITY WITH OTHER PESTICIDES

Willowood Pronamide 3.3SC is compatible with most commonly used agricultural pesticides, crop oil concentrate and adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use(s). Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

EFFECT OF SOIL TYPE, MOISTURE AND TEMPERATURE

Willowood Pronamide 3.3SC is most active in coarse to medium texture soils of low organic matter and relatively inactive in peat or muck soils or mineral soils high in

organic matter content at rates specified in the label. Herbicidal activity is best in soils containing less than 4 percent organic matter. Use in soils with higher organic matter may result in inconsistent or incomplete weed control.

The herbicidal activity of Willowood Pronamide 3.3SC is mainly through root absorption in sensitive weed species. Rain, melting snow or irrigation is essential following treatment to move Willowood Pronamide 3.3SC into the root zone of germinating weeds.

Under field conditions, Willowood Pronamide 3.3SC will remain relatively stable with little loss of herbicidal activity when soil temperatures are less than 55°F. As soil temperatures increase, degradation of the active ingredient takes place.

Willowood Pronamide 3.3SC may degrade rather quickly if left exposed on the soil surface in warm weather. If Willowood Pronamide 3.3SC is applied when air temperature exceeds 85°F, the treatment must be soil incorporated to a shallow depth (top two to three inches) or watered into the soil as soon as possible.

CULTURAL CONSIDERATIONS

For best results, apply Willowood Pronamide 3.3SC to a trash-free soil surface. Clean cultivation before application is preferable, but not necessary. To obtain optimum weed control in areas not clean cultivated, the area to be treated must be free of surface litter (dead or decaying crop and weed debris, mowing clippings, etc.). Trash-free areas create ideal conditions for rapid movement of Willowood Pronamide 3.3SC into the weed root zone following rain or irrigation.

Limitations, Restrictions, and Exceptions

TURF GROWN FOR SOD

This product may only be used on turf grown for sod or on nonresidential sites including but not limited to golf course, industrial and office building sites, stadium fields or professional athletic fields.

Product Information

Willowood Pronamide 3.3SC is a selective herbicide for the preemergence and postemergence control of annual bluegrass (*Poa annua*) from warm season grasses and the removal of perennial rye grass (*Lolium perenne*) from warm season during

spring transition. Warm season grasses include ornamental bermudagrass (*Cynodon dactylon*), Zoysiagrass, St. Augustinegrass and Centipedegrass.

Annual Bluegrass (*Poa Annua*) Control

Willowood Pronamide 3.3SC will control annual bluegrass from pre-germination and seedling stages through tillering, heading and seed formation. Willowood Pronamide 3.3SC acts slowly on seedling to mature annual bluegrass. Following application of Willowood Pronamide 3.3SC annual bluegrass may first become dark green and then gradually turn yellow and die over a 3- to 5-week period.

For effective control of annual bluegrass in turf or seed crops, moisture is necessary to move Willowood Pronamide 3.3SC in the weed root zone. Refer to the Moisture Requirements section of the label for details.

Dosage and Timing

For annual bluegrass control, Willowood Pronamide 3.3SC is applied at the rate of 1.25 to 3.5 pints of product (0.5 to 1.5 lb. active ingredient) per acre broadcast application. The dosage rate required is dependent on the growth stage of annual bluegrass at time of application.

1 One acre equals 43,560 sq. ft.

2 Use the higher rate when longer preemergence residual control is desired.

Removal of Perennial Rye Grass from Warm Season Grasses

Willowood Pronamide 3.3SC will remove postemergent perennial rye grass from warm season grasses during the spring to control the transition from cool season overseed to warm season grasses. Willowood Pronamide 3.3SC works slowly to control mature perennial rye grass. After an application of Willowood Pronamide 3.3SC, perennial rye grass will gradually die over a 4- to 6-week period. The length of this transition is dependent upon environmental factors such as temperature, rainfall and mowing height of the turf.

Dosage and Timing

For removal of perennial rye grass from warm season grasses, Willowood Pronamide 3.3SC may be applied at a rate of 1.25 to 2.5 pints of product (0.5 to 1 lb. active ingredient) per broadcast acre. It is best to apply Willowood Pronamide 3.3SC to

warm season grasses at 50% greenup. Application of Willowood Pronamide 3.3SC to dormant warm season grasses can slow greenup.

Application

Mix the specified amount of Willowood Pronamide 3.3SC in clean water and apply uniformly with a low pressure ground sprayer in 20 to 50 gallons of water per acre or 0.5 to 1 gallon of water per 1,000 sq. ft. The sprayer should be equipped with flat fan nozzles, spaced to provide uniform distribution without skips or excessive overlapping of spray patterns.

Important Note: Avoid spraying on fairways, hillsides, or approaches that may drain onto bentgrass greens or to areas overseeded with sensitive cool season grasses. Do not make an application of a wetting agent for the purpose of frost protection or soil penetration to greens or tees 14 days prior to or after a Willowood Pronamide 3.3SC application as injury may result.

Moisture Requirements

Willowood Pronamide 3.3SC acts mainly through root absorption in sensitive weed species. If no rainfall occurs within a day or so of the application, a light overhead irrigation must be made to move the chemical into the weed root zone. Avoid heavy irrigations of more than 1 inch to reduce the possibility of excess washing or leaching of the chemical from the area of application.

Willowood Pronamide 3.3SC Deactivation for Overseeding

Where it is desirable to reseed sooner than 90 days following the application of Willowood Pronamide 3.3SC, an application of an activated charcoal such as Gro-Safe, is needed. Apply the activated charcoal at the rate of 10 lbs. per 1,000 sq. ft.

Allow at least 14 days between the application of Willowood Pronamide 3.3SC and the application of charcoal for control of emerged annual bluegrass. Reseed no sooner than 7 days following charcoal application.

Turf Grown for Sod – Specific Use Restrictions

- This product may be used on non-residential seeded, sodded, or sprigged turf or turf grown for sod that is well established. Use of this product on turf that has been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress may increase the chances of turf injury.

- This product must only be applied to turf areas that are composed of the following turfgrass species:
 - o Bermudagrass (*Cynodon dactylon*)
 - o Centipedegrass (*Eremochloa ophiuroides*)
 - o St. Augustinegrass (*Stenotaphrum secundatum*)
 - o Zoysiagrass (*Zoysia japonica*)
- Avoid spraying on hill sides, fairways, or approaches that may drain onto bentgrass greens or to area overseeded with sensitive cool season grasses. Do not make an application of a wetting agent for the purpose of frost protection or soil penetration to greens or tees 14 days prior to or after a Willowood Pronamide 3.3SC application as injury may result.
- Do not apply Willowood Pronamide 3.3SC herbicide to areas that are to be overseeded with susceptible cool season grasses within 90 days of treatment unless deactivation is planned.
- Do not apply Willowood Pronamide 3.3SC to dichondra, perennial bluegrass, annual and perennial ryegrasses, fescues and bentgrasses.
- Do not graze treated areas and do not feed clippings to livestock.
- Do not apply more than 1.5 lbs./acre active ingredient (3.5 pints/acre of Willowood Pronamide 3.3SC) or make more than one application of Willowood Pronamide 3.3SC per crop season.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[Postemergence \(Weed\)](#)

[Seed forming stages](#)