

CUCUMBER, MELONS, ETC. - PICKLEWORM, MELONWORM, ETC.

General Information

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children or pets to enter the treated area until dusts have settled. Do not apply to trees.

To control certain insects and to control or prevent certain blight, leaf spots, mildew and certain other diseases on listed plants. Use as a dust any time conditions are favorable, as soon as plants are above the ground or as soon as insects appear. Keep foliage covered, especially new growth.

When dusting, wear household latex or rubber gloves. Apply only from this shaker bottle. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin. For plants above chest height consider using a liquid carbaryl product.

Start dusting at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before re-entering.

AS A DUST

This bottle is a squeeze type duster. Shake bottle to fluff powder. Press sides of bottle with fingers, using rapid motion for fog effect.

WHEN TO APPLY

For insects apply when insects appear. Young insects and larvae are usually easier to control than adults. When blossoms are in full bloom, protect pollinators by using a fungicide only, such as Bonide Copper Dust. For disease, applications should begin well before they are expected to appear.

Limitations, Restrictions, and Exceptions

Apply every 5 days; do not exceed 8 applications per season.

Method

[Broadcast/Foliar Air](#)

Broadcast/Foliar Ground

Pre-Harvest Interval

3 days

Rates

field rates 0

•

Timings

When insects appear.