

## **CEREAL GRAINS - ARMYWORM, BEAN LEAF BEETLE, ETC.**

### General Information

#### GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gal per acre by air or 10 gal per acre by ground unless otherwise specified in the label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Lamdec Insecticide may be applied before, during or after planting. For soil incorporated applications, use higher rates for improved control.

#### RESISTANCE MANAGEMENT

Lamdec Insecticide is a Group 3 Insecticide (contains the active ingredient lambdacyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Refer the product label for the Rate Conversion Chart.

#### Limitations, Restrictions, and Exceptions

## CEREAL GRAINS

### Pests:

Armyworm, Fall Armyworm, Yellowstriped Armyworm - Use higher rates for large larvae.

Bird Cherry-Oat Aphid, Corn Leaf Aphid, English Grain Aphid - Suppression only.

European Corn Borer, Hop Vine Borer, Southwestern Corn Borer, Stalk Borer - For control before the larva bores into the plant stalk or ear.

Tobacco Budworm - For control before the larva bores into the plant stalk or ear. See Resistance statement under General Directions for Use.

### Remarks:

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gal of water per acre.
- Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment.
- Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- Do not apply more than 0.06 lb ai (7.68 fl oz or 0.48 pt of product) per acre after silk initiation.
- Do not apply more than 0.03 lb ai (3.84 fl oz or 0.24 pt of product) per acre after corn has reached the milk stage (yellow kernels with milky fluid).

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

[field\\_rates 2](#)

•

Restricted Entry Interval

24 hours

Timings

[Upon insect populations reaching locally determined economic threshold.](#)

[European Corn Borer, Hop Vine Borer, Southwestern Corn Borer, Stalk Borer - Before the larva bores into the plant stalk or ear.](#)