

GRAPES

General Information

GENERAL INSTRUCTIONS

The per acre use rates for NORDOX 30/30 WG are applicable for dilute, concentrate and aerial sprays. Where dosage and/or interval ranges are indicated, these may be adjusted according to disease conditions. For example, when conditions favor severe disease pressure, use the maximum rate and shortest spray interval. High rates may be phytotoxic when crop is experiencing environmental stress (hot, dry conditions).

FROST INJURY PROTECTION

Application of NORDOX 30/30 WG made to all crops listed on the label at rates indicated on the label, just prior to anticipated frost conditions, will sustain control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide protection against light frost. Not intended for those geographic areas where weather conditions favor severe frost.

Refer to the product label for the minimum spray volume (gallons) per acre.

Limitations, Restrictions, and Exceptions

GRAPES

- Apply just before bud break, when the shoots are 6-8 inches long, just after bloom, and throughout season as needed. Minimum retreatment interval is 3 days. Foliar injury may occur on coppersensitive varieties.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

Just before bud break, when the shoots are 6-8 inches long, just after bloom.