

WHEAT, BARLEY, OATS, AND RYE - PERENNIAL WEEDS

General Information

USE PRECAUTIONS

Unless otherwise noted under individual DIRECTIONS section, for aerial application, apply the recommended amount in a minimum of 2 gallons of water per acre. For ground application, apply the recommended amount in a minimum of 10 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. DO NOT apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift. Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

WHEAT, BARLEY, OATS, AND RYE

DIRECTIONS

Apply when weeds are small.

Apply after crop has reached the 3 to 5 leaf stage up to early boot stage.

For less susceptible weeds, use the 1 5/8 pint rate after the crop has tillered and up to early boot stage. Do not spray from boot stage to dough stage.

- The higher rate may cause grain injury. Any legumes present may be killed.

RESTRICTIONS AND LIMITATIONS FOR SMALL GRAINS

Do not forage or graze treated areas for 7 days after application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

Broadcast/Foliar Ground

Rates

field rates 0

field rates 1

-

Restricted Entry Interval

12 hours

Timings

When weeds are small.

After crop has reached the 3 to 5 leaf stage up to early boot stage.