

PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES - CANADA THISTLE AND WHITETOP (HOARY CRESS)

General Information

USE PRECAUTIONS

Unless otherwise noted under individual DIRECTIONS section, for aerial application, apply the recommended amount in a minimum of 2 gallons of water per acre. For ground application, apply the recommended amount in a minimum of 10 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. DO NOT apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift. Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES

DIRECTIONS

Use higher rate for Whitetop, Canada thistle, and other less susceptible weeds. Spray perennial in early bud to full bloom stage and during active regrowth in the fall.

- Do not use on alfalfa, clover, other legumes or newly seeded areas.

RESTRICTIONS AND LIMITATIONS FOR PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAMS AND SET-ASIDE ACRES

Do not forage or graze treated areas for 7 days after application.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[To established grasses before the head comes into boot stage and on seedling grasses after the grass has at least 5 leaves \(tillered \).](#)