

NON-CROP AREAS - SPOT TREATMENT (CANADA THISTLE, WHITETOP (HOARY CRESS) AND BUTTERCUP)

General Information

USE PRECAUTIONS

Unless otherwise noted under individual DIRECTIONS section, for aerial application, apply the recommended amount in a minimum of 2 gallons of water per acre. For ground application, apply the recommended amount in a minimum of 10 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. DO NOT apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift. Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

NON-CROP AREAS

DIRECTIONS

Spray when weeds are in bud to early bloom and again during active fall regrowth.

Apply in 3 to 4 gallons of water. Spray to thoroughly wet weeds.

RESTRICTIONS AND LIMITATIONS FOR NON-CROP AREAS

Do not forage or graze treated areas for 7 days after application.r.

Method

[Spot treatment](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

When weeds are in bud to early bloom and again during active Fall regrowth.