

# **ALMONDS, APRICOTS, CHERRY, ETC. - BROWN MITE, EUROPEAN MITE, ETC.**

## General Information

### GROUND EQUIPMENT APPLICATIONS

Release Height: Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

Wind Speed: Do not apply at wind speeds greater than 15 mph at the application site.

Droplet Size: Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

### AIRBLAST APPLICATIONS

Turn off outward pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

### USE LIMITATIONS AND PRECAUTIONS

Preharvest Interval - This product may be used up to the day of harvest. To ensure thorough coverage, dilute applications (greater than 150 gallons of spray solution per acre) should be used. Concentrate applications (usually from 20.0 to 150 gallons of spray solution) may reduce the coverage and thus the effectiveness. Application volumes depend on the crop type, crop size and target pest. Adjust the spray volume to obtain the best coverage without runoff. Extreme care should be taken when using concentrate sprays as the potential for enhanced crop phytotoxicity is increased. A concentrate application can provide satisfactory results as long as the spray unit is properly engineered, calibrated and operated.

Note: Use efficient equipment of the proper type. Avoid spraying during or immediately prior to hot or freezing weather (over 95 °F or under 32 °F), hot dry winds, rain or other unsuitable conditions; avoid over spraying or double spraying. Plants should be sprayed only when in vigorous condition and when their moisture

condition is suitable. Before using, make certain spray tank is free of sulfur residues.

Do not apply sulfur within 30 days except in the Northwest (Oregon and Washington) where it may be used in combination on Pears as a Post Harvest (after old fruit is off the tree), Dormant and Delayed Dormant (before the scales slip, or at or prior to bud swell) treatment and on Apples prior to or up to the Delayed Dormant (1/2 inch green) period. In areas west of the Continental Divide do not apply to captan treated trees within 90 days after or 60 days before this oil. In areas east of the Continental Divide do not apply to captan treated trees within 10 to 14 days before and after this oil. If in doubt, spray a portion of one tree previously treated with sulfur or captan using 1.0 quart to 25.0 gallons water. Burn will show in several days if sufficient time has not elapsed.

This product has not been tested on all species or varieties. Before treating a large area, treat a small area and observe prior to full scale application. A period of 14 days must elapse before any of the following chemicals are used before or after an application of this product: anilazine (Dyrene), chlorothalonil (Bravo), dicloran (Botran), dicofol (Kelthane), dinocap (Dikar, Karathane), permethrin (Ambush, Pounce). Do not use chlorothalonil (Bravo) or dimethoate (Cygon) in a spray program with this product on Grapes, Ornamentals and Strawberries. Do not use dicofol (Kelthane) in a spray program with this product on Ornamentals and Strawberries. Do not use propargite (Omite) with an oil spray or within 30 days before or after an oil spray.

#### Limitations, Restrictions, and Exceptions

ALMONDS, APRICOTS, CHERRY, ETC. - BROWN MITE, EUROPEAN MITE, ETC

#### Rate:

Concentrate Spray: Do not use more than 4.0 gal/A in a minimum of 20.0 gal water/A.

#### Comments:

Certain varieties of Plums and Prunes may be injured by oil sprays, check for tolerance prior to treatment.

Do not apply to trees lacking moisture.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

- 

Restricted Entry Interval

4 hours

Timings

[N. A.](#)