

# **CONTROL OF MICE**

## General Information

READ THIS LABEL and follow all use directions and precautions.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents: 1. Store product not in use in a location out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations.

These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** This product may be used to control Norway rats, roof rats and house mice at the Urban and Non-urban sites identified below. Do not place bait where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**Urban Areas:** This product may only be used: in and around homes; in and around industrial, commercial and public buildings; in transport vehicles (ships, trains, aircraft), and in and around related port or terminal buildings. All applications "around" buildings must be made in tamper-resistant bait stations placed along the walls or other external parts of buildings (e.g., doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

**Non-urban Areas:** This product may only be used inside of homes and agricultural buildings.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice are likely to find and consume bait, such as along walls,

by gnawed openings, in corners and concealed places, between floors or walls, or locations where rodents or their signs are seen. Remove as much alternative food as possible.

Follow-up: Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

#### Limitations, Restrictions, and Exceptions

Each Generation Block is scored to be broken into 2-ounce pieces

Mice: Break block and apply one 2-ounce piece per placement.

Space placements at 8 to 12 foot intervals. Maintain a constant supply of fresh bait for 15 days or until signs of mouse activity cease.

Follow-up: Replace contaminated or spoiled bait immediately.

Collect and dispose of all visible dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

#### Method

##### [Bait](#)

##### Timings

[Until signs of house mouse activity cease.](#)