

RANGELAND - USDA APHIS PROGRAM

General Information

For the control of cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, sowbugs and ants (except fire ants, harvester ants, pharoah ants and carpenter ants).

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl residues when Carbaryl is used in accordance with label directions.

If Carbaryl insecticide is used in accordance with label directions, carrot tops, sugar beet tops and corn forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

APPLICATION METHOD

This product may be applied only by ground using a mechanical spreader or underground shank applicator. Do not apply by hand, spoon or belly grinder. EXCEPTION: Aerial applications only to rangeland are allowed only under the auspices of the Animal and Plant Health Inspection Service (APHIS) Rangeland Grasshopper and Mormon Cricket Suppression Program, and/or allied state programs.

Limitations, Restrictions, and Exceptions

USE RATE

USDA APHIS PROGRAM for Mormon Cricket and Grasshoper Suppression

Apply 4 lbs. per acre per application with a maximum yearly application rate of 8 lbs. per acre.

- Retreatment interval = 14 days

- Do not apply within 14 days of grazing
- Use of a flaky cereal grain bait is prohibited.

Aerial application is allowed only through USDA Aphis and affiliated state grasshopper and Mormon Cricket suppression programs.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[N.A.](#)