

CORN - PREPLANT AND BURNDOWN

General Information

USE INFORMATION

ET herbicide/defoliant is designed for use as a contact herbicide for broadleaf weed control, defoliation, and desiccation.

For best results, use ET herbicide/defoliant for control of annual or perennial herbaceous broadleaf weeds less than 4 inches in height, or rosettes less than 3 inches in diameter. Use the higher rates and spray volumes for control of larger weeds; control may be reduced with weeds larger than 4 inches.

ET herbicide/defoliant must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

Use an approved agricultural buffering agent, buffering to less than pH 7.5, if using ET herbicide/defoliant in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding ET herbicide/defoliant to the spray tank.

ET herbicide/defoliant is a contact herbicide and defoliant and requires thorough coverage for complete broadleaf weed control and defoliation/desiccation.

See label for Rational Crop Restrictions

Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground unless otherwise specified.

Do not apply ET herbicide/defoliant through any type of irrigation system.

ET herbicide/defoliant is rainfast within one hour after application.

See Supplemental label for Rotational Crop Restriction

See Supplemental label For Use on Triticale

Limitations, Restrictions, and Exceptions

CORN

- Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground.
- Do not apply more than 2.0 fl oz/acre per season prior to planting and/or emergence of crop.
- Allow a minimum of 30 days between applications for this use.
- The addition of a spray tank adjuvant at a concentration of 0.5% to 2.0% is recommended for optimum weed control.
- Refer to page 8 for crop rotations/plantback restrictions.
- Use the higher rate for hard-to-control weeds such as field bindweed and kochia.

Corn – All Uses: Do not apply more than 3.0 fl oz/acre per growing season to corn.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[At-Plant](#)

[Preemergence \(Crop\)](#)

Preplant