

TRITICALE AND WHEAT - PREPLANT AND BURNDOWN

General Information

USE INFORMATION

ET herbicide/defoliant is designed for use as a contact herbicide for broadleaf weed control, defoliation, and desiccation.

For best results, use ET herbicide/defoliant for control of annual or perennial herbaceous broadleaf weeds less than 4 inches in height, or rosettes less than 3 inches in diameter. Use the higher rates and spray volumes for control of larger weeds; control may be reduced with weeds larger than 4 inches.

ET herbicide/defoliant must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

Use an approved agricultural buffering agent, buffering to less than pH 7.5, if using ET herbicide/defoliant in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding ET herbicide/defoliant to the spray tank.

ET herbicide/defoliant is a contact herbicide and defoliant and requires thorough coverage for complete broadleaf weed control and defoliation/desiccation.

See label for Rational Crop Restrictions

Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground unless otherwise specified.

Do not apply ET herbicide/defoliant through any type of irrigation system.

ET herbicide/defoliant is rainfast within one hour after application.

See Supplemental label for Rotational Crop Restriction

See Supplemental label For Use on Triticale

Limitations, Restrictions, and Exceptions

SOYBEAN

- Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground.
- ET herbicide/defoliant can be applied from crop emergence to the V6 growth stage.
- Allow a minimum of 30 days between applications for this use.
- Do not use crop oils or crop oil concentrates for postemergence applications.
- Do not graze soybean forage or cut for hay within 7 days of last ET herbicide/defoliant application.
- Some temporary herbicidal leaf speckling may appear on the crop. This effect is transient and will NOT appear on new growth.
- Refer to page 8 for crop rotations/plantback restrictions.
- Use the higher rate for hard-to-control weeds such as field bindweed and kochia.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Burndown](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[At-Plant](#)

[Preemergence \(Crop\)](#)

[Preplant](#)