

GUAYULE

General Information

USE INFORMATION

ET herbicide/defoliant is designed for use as a contact herbicide for broadleaf weed control, defoliation, and desiccation.

For best results, use ET herbicide/defoliant for control of annual or perennial herbaceous broadleaf weeds less than 4 inches in height, or rosettes less than 3 inches in diameter. Use the higher rates and spray volumes for control of larger weeds; control may be reduced with weeds larger than 4 inches.

ET herbicide/defoliant must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

Use an approved agricultural buffering agent, buffering to less than pH 7.5, if using ET herbicide/defoliant in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding ET herbicide/defoliant to the spray tank.

ET herbicide/defoliant is a contact herbicide and defoliant and requires thorough coverage for complete broadleaf weed control and defoliation/desiccation.

See label for Rational Crop Restrictions

Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground unless otherwise specified.

Do not apply ET herbicide/defoliant through any type of irrigation system.

ET herbicide/defoliant is rainfast within one hour after application.

See Supplemental label for Rotational Crop Restriction

See Supplemental label For Use on Triticale

Limitations, Restrictions, and Exceptions

GUAYULE

Use Directions and Restrictions

- Allow a minimum of 30 days between applications.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- NO portion of the treated field, including seed, seed screenings, or stubble, may be used for human consumption.
- This information must be passed along to anyone using this crop.
- Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

For best results, use ET herbicide for control of annual broadleaf weeds when weeds are less than 4" in height or rosettes are less than 3" in diameter. ET is a contact herbicide. Thorough uniform spray coverage is essential for good control. Phytotoxicity may occur as leaf speckling; however, no crop loss is expected. Addition of a crop oil concentrate (COC) or nonionic surfactant is recommended for optimum control. Oil base spreaders will improve weed control but may also increase crop injury. Use with caution. Follow manufacturer's recommended use rates. Read and follow all label directions for each tank mix product. Always use in accordance with the most restrictive of label precautions and limitations.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[When weeds are less than 4" in height or rosettes are less than 3" in diameter.](#)