

CORN (FIELD CORN, ETC.)- POSTEMERGENCE

General Information

USE INFORMATION

ET herbicide/defoliant is designed for use as a contact herbicide for broadleaf weed control, defoliation, and desiccation.

For best results, use ET herbicide/defoliant for control of annual or perennial herbaceous broadleaf weeds less than 4 inches in height, or rosettes less than 3 inches in diameter. Use the higher rates and spray volumes for control of larger weeds; control may be reduced with weeds larger than 4 inches.

ET herbicide/defoliant must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

Use an approved agricultural buffering agent, buffering to less than pH 7.5, if using ET herbicide/defoliant in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding ET herbicide/defoliant to the spray tank.

ET herbicide/defoliant is a contact herbicide and defoliant and requires thorough coverage for complete broadleaf weed control and defoliation/desiccation.

See label for Rational Crop Restrictions

Apply ET herbicide/defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground unless otherwise specified.

Do not apply ET herbicide/defoliant through any type of irrigation system.

ET herbicide/defoliant is rainfast within one hour after application.

See Supplemental label for Rotationcal Crop Restriction

See Supplemental label For Use on Triticale

Limitations, Restrictions, and Exceptions

CORN

- Allow a minimum of 30 days between applications for this use.
- Use the higher rate for more mature plants. Control will be reduced when Volunteer Cotton is greater than the 4-leaf stage
- Apply ET Herbicide/Defoliant in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground.
- ET Herbicide/Defoliant can be applied from crop emergence to the V2 growth stage.
- Do not apply postemergence to sweet corn.
- Do not make more than 1 application per season for this use.
- Do not apply more than 0.75 fl oz/acre per season for this use.
- Do not use crop oils or crop oil concentrates for postemergence applications.
- Some temporary herbicidal leaf speckling may appear on the crop. This effect is transient and will NOT appear on new growth.
- Do not harvest corn for silage within 50 days after last application of ET Herbicide/Defoliant.
- Do not harvest corn for grain or stover within 90 days after last application of ET Herbicide/Defoliant

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

Postemergence (Crop)