

CUT STUBBLE

General Information

GENERAL INFORMATION

Use Sites: Arsenal is an aqueous solution to be mixed with water and a surfactant and applied as a spray solution to grass pasture and rangeland and noncropland areas such as railroad, utility, pipeline and highway rights-of-way, utility plant sites, petroleum tank farms, pumping installations, fence rows, storage areas, non-irrigation ditchbanks, including grazed or hayed areas within these sites.

Arsenal is recommended for the establishment and maintenance of wildlife openings. Arsenal may also be used for the release of unimproved bermudagrass (see specific directions) and for use under certain paved surfaces (see specific directions).

Application Methods: Arsenal will control most annual and perennial grasses and broadleaf weeds in addition to many brush and vine species and Arsenal will provide residual control of labeled weeds which germinate in the treated areas. This product may be applied either preemergence or postemergence to the weeds; however, postemergence application is the method of choice in most situations, particularly for perennial species. For maximum activity, weeds should be growing vigorously at the time of postemergence application and the spray solution should include a surfactant (See Adjuvant Section for specific recommendations). These solutions may be applied selectively by using low-volume techniques or may be applied broadcast by using ground equipment or aerial equipment. In addition, Arsenal may also be used for stump and cut stem treatments (see specific directions).

Herbicidal Activity: Arsenal is readily absorbed through leaves, stems, and roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into, and kills, underground storage organs which prevents regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two weeks after application.

Complete kill of plants may not occur for several weeks. Applications of Arsenal herbicide are rainfast one hour after treatment.

PRECAUTIONS FOR AVOIDING INJURY

TO NON-TARGET PLANTS

Untreated trees can occasionally be affected by root uptake of Arsenal through movement into the top soil.

Injury or loss of desirable trees or other plants may result if Arsenal is applied on or near desirable trees or other plants, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots.

Aerial Application Methods and Equipment: Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

Ground Application (Broadcast): Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

BRUSH CONTROL

AERIAL APPLICATIONS:

All precautions should be taken to minimize or eliminate spray drift. Fixed wing aircraft and helicopters can be used to apply Arsenal, however, DO NOT make

applications by fixed wing aircraft unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or, when treating open tracts of land, spray drift as a result of fixed wing aircraft application can be tolerated. Aerial equipment designed to minimize spray drift, such as a helicopter equipped with a Microfoil boom, Thru-Valve boom or raindrop nozzles, must be used and calibrated.

Except when applying with a Microfoil boom, a drift control agent may be added at the recommended label rate. To avoid drift, applications should not be made during inversion conditions, when winds are gusty, or any other conditions which allow drift. Side trimming is not recommended with Arsenal unless death of treated tree can be tolerated.

Uniformly apply the recommended amount of Arsenal in 5 to 30 gallons of water per acre; include in the spray solution a nonionic surfactant or methylated seed oil or manufacturer's label rate of a silicone-based surfactant (See the Adjuvant section of the label for specific recommendations).

A foam reducing agent may be added at the recommended label rate, if needed.

IMPORTANT: Thoroughly clean application equipment, including landing gear, immediately after use of this product. Prolonged exposure of this product to uncoated steel (except stainless steel) surfaces may result in corrosion and failure of the exposed part. The maintenance of an organic coating (paint) may prevent corrosion.

GROUND APPLICATIONS:

IMPORTANT: To minimize spray drift, select proper nozzles to avoid spraying a fine mist, use pressures less than 50 psi, and do not spray under gusty or windy conditions.

Add a foam reducing agent, if needed, and a spray pattern indicator, if desired, at the recommended label rates. Clean application equipment after using this product by thoroughly flushing with water.

When making applications to rights-of-way corridors where desirable tree roots may extend, use 1 to 3 pints of Arsenal herbicide per acre in combination with recommended tank mixes. It is not recommended to use rates higher than 3 pints

per acre in these situations as injury or death of desirable trees may occur when their roots extend into treated zones.

Side Trimming:

DO NOT side trim with Arsenal unless severe injury or death of the treated tree can be tolerated. Arsenal is readily translocated and can result in death of the entire tree.

Low Volume:

Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. To prepare the spray solution, thoroughly mix in water 0.5 to 5% Arsenal plus surfactant (See the Adjuvant section of the label for specific recommendations). A foam reducing agent may be applied at the recommended label rate, if needed. For control of difficult brush species (see Weeds Controlled section for relative susceptibility of weed species), use the higher concentrations of herbicide and/or spray volumes but do not apply more than 6 pints of Arsenal per acre. Excessive wetting of foliage is not recommended. See the Mixing Guide below for some suggested volumes of Arsenal and water.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

CUT STUBBLE

Arsenal can be applied within 2 weeks after mechanical mowing or cutting of brush. To suppress or control resprouting, uniformly apply a spray solution of Arsenal at the rate of 1 to 2 pints per acre to the cut area. Arsenal may be tank mixed with Tordon K or picloram to aid in control or suppression of brush. The addition of 5% (v/v) or more of a penetrating agent can aid in uptake through the bark or exposed roots.

Cut stubble applications are made to the soil and cut brush stumps. This type of application may increase ground cover injury. However, vegetation will recover.

Making applications of Arsenal directly to the soil can increase potential root uptake causing injury or death of desirable trees.

Efficacy can be increased and root uptake by desirable vegetation can be decreased

if the brush is allowed to regrow and the foliage is treated. See the Brush Control section of the label.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[2 weeks after mechanical mowing or cutting of brush.](#)