

## **CORN, FIELD AND SWEET**

### General Information

#### GENERAL RECOMMENDATIONS

B-17 ACID SPRAY at a rate of 5.9 lbs. provides the equivalent of 1 lb. of boron. B-17 contains no sodium and is mildly acidic.

DILUTION: Apply the suggested rates of B-17 BORIC ACID SPRAY in the following amounts of water per acre. Minimum dilution rate - 1 lb. per 5 gallons of water per suggested rate.

#### Field & Row Crops

By air: 10-20 gals.

By ground: 10 gals. minimum.

#### Tree Crops

By air: 10 gals. minimum.

By ground concentrate: 40-100 gals.

By ground dilute: 100-500 gals.

In oil sprays, use 1-5 lbs. per 100 gallons.

COMPATIBILITY: This product has been designed to be compatible with most commonly used insecticides, fungicides and nutritional products. However, a compatibility test is recommended for combinations that have not been used previously. Consult your NUTRI-TECH representative for detailed compatibility information.

FOR ALL CROPS- DO NOT EXCEED THE RATE OF 6.5 LBS. PER ACRE PER YEAR OF B-17 IN FOLIAR SPRAYS, 12 LBS. PER ACRE IN SOIL APPLICATIONS. SUGGESTED RATES MAY BE APPLIED IN SPLIT APPLICATIONS.

#### Limitations, Restrictions, and Exceptions

## CORN, FIELD AND SWEET

- Apply 0.5-3 lbs. when 10-20" high, repeat in 2-4 weeks as necessary

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Rates

[field\\_rates 0](#)

- 

### Timings

[When 10-20\" high.](#)