

# **TREE CROPS - PEARS (POSTHARVEST, DORMANT THROUGH PRE-BLOOM)**

## General Information

### GENERAL INSTRUCTIONS

Apply AMBUSH Insecticide as shown in Directions for Application. Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. Do not apply more than 2.0 lbs. a.i./A per season. Crops in the label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.

### Resistance

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

### Limitations, Restrictions, and Exceptions

### Remarks

Apply by air or ground equipment. By air, apply in a minimum of 10 gal. of water/A. By ground, apply in a minimum of 25 gal. of water/A. Use sufficient water to obtain uniform coverage. Apply at a rate of 0.8-1.6 pts. (0.2 to 0.4 lbs. a.i.)/A. May be combined with 2-8 gal. of spray oil/A. Apply during the postharvest through dormant

through pre-bloom growth periods only. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.

Pre-bloom: Apply 2 prebloom sprays. Pre-bloom sprays can be applied from dormant through bud burst stages.

Post-harvest: For control of Codling Moth: Apply 0.2-0.4 lbs. a.i./A (12.8-25.6 fl. oz./A). AMBUSH Insecticide may be applied with or without dormant oil by ground equipment (dilute or concentrate sprays) or by aircraft. When combined with dormant oil and applied by air, observe the application rate and spray volume specified on the dormant oil label. When not using dormant oil, pears may be treated by air with a minimum of 15 gal. of finished spray/A. Apply by air only when soil conditions do not permit regular ground application. For maximum effectiveness, combine AMBUSH Insecticide with labeled rates of dormant oil and apply with air blast ground sprayers in volumes of 100-400 gal. of finished spray/A.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

[field rates 3](#)

•

#### Restricted Entry Interval

12 hours

#### Timings

[Post-harvest](#)

[Dormant through Pre-bloom](#)