

# **LEAF-MAX APPLICATION**

## General Information

### GENERAL RECOMMENDATIONS

Leaf-MAX II is designed for foliar use to help maximize nutritional requirements needed for proper leaf development, plant growth, and bud quality. It is most effective when applied just prior and during strong vegetative growth phases. Can also be used post-harvest to help nutrient storage and bud strength. When applying post-harvest, leaves must be in good condition for maximum benefit. Do not apply on plants where leaf senescence has begun, as benefits will be limited.

### Limitations, Restrictions, and Exceptions

### IRRIGATION STATEMENT

Applications can be made through certain irrigation systems including, but not limited to, low volume systems and typical agricultural sprinkler systems. The specific injection equipment and irrigation system must conform to all federal, state and local regulations pertaining to the fertilization of plants through an irrigation system. Accurate and properly calibrated injection equipment and an irrigation system that is operated at maximum distribution uniformity will help avoid plant injury and/or product ineffectiveness. For best results, with the exception of center pivot and lateral systems, apply this product toward the end of the irrigation set. Allow sufficient time for the product to be flushed completely out of the entire irrigation system prior to turning off the irrigation water. Contact your Certified Crop Advisor for specific recommendations for your area.

### Method

[Foliar spray](#)

[Irrigation](#)

[Foliar spray](#)

[Irrigation](#)

[Foliar spray](#)

[Irrigation](#)

Rates

field rates 0

- 

Timings

At the beginning of and during a strong foliar growth.

When tissue analysis shows a need for zinc and/or magnesium,.

Within 2-4 weeks post harvest.