

HIGHWAY, ROADSIDE, PIPELINE AND UTILITY RIGHTS-OF-WAY, INDUSTRIAL AREAS, FENCE ROWS AND OTHER NON-CROP SITES

General Information

RESISTANCE MANAGEMENT

Some weeds are known to develop resistance to herbicides that have been used repeatedly. While the development of herbicide resistance is well understood, it is not easily predicted. Therefore herbicides should be used in conjunction with the resistance management strategies in the area. Consult the local or State agricultural advisors for details. If herbicide resistance should develop in the area, this product used alone may not continue to provide sufficient levels of weed control. If the reduced levels of control can not be attributed to improper application techniques, improper use rates, improper application timing, unfavorable weather conditions or abnormally high weed pressure, a resistant strain of weeds may have developed.

To reduce the potential for weed resistance use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the recommended rates and in accordance with the use directions. Do not use less than recommended label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when weeds are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

CALIFORNIA ONLY SPECIFIC RESTRICTIONS ON APPLICATIONS OF Portfolio 4F CA.

Artificial Recharge Basins. Do not use below the high water line inside artificial recharge basins (a surface facility, such as an infiltration pond or basin, or spreading ground that is specifically designed and managed to increase the infiltration of introduced surface water supplies into a ground water basin), unless this product is applied six months or more before the basin is used to recharge ground water.

Unlined Canals and Ditches. Do not use below the high water line inside unlined canals and ditches unless either (a) the pesticide user can document that the

percolation rate of the canal or ditch is equal to or less than 0.2 inches per hour (0.002 gallons per minute per square foot), or (b) the pesticide is applied six months before water is run in the canal or ditch.

Rights-of-Way. Do not use on engineered rights-of-way in areas established by the California Department of Pesticide Regulation as leaching or runoff ground water protection areas* unless either (a) any runoff from the treated right-of-way shall pass through a noncrop fully vegetated area adjacent, and equal in area, to the treated area, or spread out onto an adjacent unenclosed fallow field that is at least 300 feet long and that will not be irrigated for six months following application with the exception of the addition of adequate moisture that is required for herbicidal activation following application as described under General Application Instructions, with full consideration of any plantback restrictions, or (b) the property operator complies with any permit issued pursuant to the storm water provisions of the federal Clean Water Act pertaining to the treated area.

Runoff Groundwater Protection Areas. Do not use in areas identified by the California Department of Pesticide Regulation as a runoff ground water protection area unless one of the following management practices can be met: (a) Soil disturbance. Within seven days before this product is applied, the soil to be treated shall be disturbed by using a disc, harrow, rotary tiller, or other mechanical method. This subsection does not apply to the area to be treated that is immediately adjacent to the crop row and that do not exceed 33 percent of the distance between crop rows or, in citrus, to the band from the tree row to the dripline; or

(b) Incorporation of the pesticide. Within 48 hours after the day this product is applied, the pesticide shall be incorporated on at least 90 percent of the area treated; using a disc, harrow, rotary tiller, or other mechanical method, or by sprinkler or low flow irrigation, including chemigation where allowed by the label, using a minimum of 1/4 inch of irrigation water and a maximum of one inch as described under General Application instructions at application rates that do not cause surface water runoff from the treated property or to wells on the treated property; or

(c) Band treatment. This product is applied as a band treatment immediately adjacent to the crop row so that not more than 33 percent of the distance between rows is treated or, in citrus, not more than the area from the tree row to the dripline is treated; or

(d) Timing of application. This product is applied between April 1 and July 31; or

(e) Retention of runoff on field. For six months following the application, the field shall be designed, by berms, levees, non-draining circulation systems, to retain all irrigation runoff and all precipitation on, and drainage through, the field. The retention area on the field shall not have a percolation rate of more than 0.2 inches per hour (5 inches per 24 hours); or

(f) Retention of runoff in a holding area off the field. For six months following the application, all runoff shall be channeled to a holding area off the application site, under the control of the property operator, that is designed to retain all irrigation runoff and all precipitation on, and drainage through, the treated field and all other areas draining into that holding area. The holding area shall not have a percolation rate of more than 0.2 inches per hour (5 inches per 24 hours); or

(g) Runoff onto a fallow field. For six months following application, runoff shall be managed so that it runs off onto an adjacent unenclosed fallow field at least 300 feet long that is not irrigated for six months after application with the exception of the addition of adequate moisture that is required for herbicidal activation following application as described under General Application Instructions, with full consideration of any plant back restrictions.

Leaching Ground Water Protection Areas. Do not use in areas designated by the California Department of Pesticide Regulation as leaching ground water protection areas* unless either (a) the user does not apply any irrigation water for six months following application of this product or (b) the user applies this product to the planting bed or the berm above the level of irrigation water in the furrow or basin and the water level shall remain at or below that level for six months following application of the pesticide with the exception of the addition of adequate moisture that is required for herbicidal activation following application as described under General Application Instructions, or (c) irrigation is managed so that the ratio of the amount of irrigation water applied divided by the net irrigation requirement is 1.25 or less for six months following application of this product.

* Consult with your County Agricultural Commissioner to determine whether the application will be within an area designated by the California Department of Pesticide Regulation as either a Runoff Ground Water Protection Area or a Leaching Ground Water Protection Area. Details regarding the locations of these Areas are also available via the internet at www.cdpr.ca.gov/docs/emon/grndwtr/gwp_regs.htm

Highway, Roadside, Pipeline and Utility Rights-of-Way, Industrial Areas, Fence Rows, and Other Non-crop Sites.

GENERAL INFORMATION

Portfolio 4F CA is a selective soil applied herbicide for the control of certain broadleaf weeds, grasses and sedges. When applied according to directions, it will provide control of susceptible species. Portfolio 4F CA is formulated as flowable (suspension concentrate) containing four pounds of the active ingredient sulfentrazone per gallon.

The mode of action of Portfolio 4F CA involves uptake by weed roots and shoots. Observe all instructions, mixing directions, application precautions and other label information of each product when tank mixing with Portfolio 4F CA.

Proper handling instructions: This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned

wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well, are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

GENERAL APPLICATION INFORMATION

Utilize a boomless application system or a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Utilize nozzles and boomless sprayer configurations which produce minimal amounts of fine spray droplets. Do not exceed 25 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles or boomless application systems. Apply a minimum of 10 gallons of finished spray per acre.

Water must be used as the carrier for this product when applied alone, or when tank mixed with other herbicides.

Avoid letting this product sit overnight as settling of product and difficulty of

resuspending may occur.

Do not allow spray to drift onto adjacent plants as injury to other plants may occur.

Do not apply to ornamental shrubs and trees, turf grasses or crops.

Limitations, Restrictions, and Exceptions

APPLICATION INSTRUCTIONS

Highway, Roadside, Pipeline and Utility Rights-of-Way.

Portfolio 4F CA can be used to control many weeds and maintain bare ground in highway, roadside, pipeline and utility rights-of-way. Such areas would include, but are not limited to, guard rails, road shoulders, electric utility substations, pipeline pumping stations, around electric transmission towers, around distribution line poles and in other areas where complete vegetation control is desired.

Industrial Areas, Fence Rows and Other Non-crop Sites.

Portfolio 4F CA controls weeds and maintains bare ground in industrial areas including production facilities, tank farms, storage areas, parking areas, lumber yards, airports, military installations, along fence rows, and in similar non-crop sites where complete vegetation control is needed.

Method and Rate of Application

For residual control of germinating weeds in non-crop land, apply this product as a broadcast treatment at 8 to 12 fluid ounces (0.25 to 0.375 pounds active ingredient) per acre by ground in a minimum of 10 gallons of spray solution per acre.

DO NOT apply Portfolio 4F CA to soils classified as sand with less than 1% Organic Matter.

Use labeled rates of burndown herbicides such as glyphosate, glyphosate - trimesium, diquat, 2,4-D, dicamba, etc. as tank mixtures with Portfolio 4F CA. Use recommended adjuvants for the herbicide tank mix partner. For all products used in tank mixes, refer to the specific product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

Timing

For best results, apply Portfolio 4F CA alone or in combination with other herbicides for residual control of weeds in late summer, fall, or early spring to insure adequate moisture for soil activation.

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[In late summer, fall, or early spring.](#)