

BROADLEAF (DICOT) WEEDS

General Information

Application Timing: The best weed control is obtained when NEGATE 37WG is applied to young, actively growing weeds. The degree and duration of control may depend on (a) weed spectrum and infestation intensity, (b) weed size at application, and (c) environmental conditions at and following treatment.

Precaution: Tolerance of the majority of cultivars of Zoysiagrass to NEGATE 37WG has not been fully investigated. It is known that Emerald, Zenith, and Meyer cultivars of Zoysiagrass have shown tolerance to similar to that of bermudagrass. The effects of NEGATE 37WG on these turfgrasses during transition have not been fully evaluated.

IMPORTANT Addition of a nonionic surfactant of at least 80% active ingredient at 0.25 percent by volume (1 qt./100 gal) provides maximum performance, but may temporarily increase chlorosis of the turf.

When an adjuvant is to be used with this product, Control Solutions, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant. Products must contain only EPA-exempt ingredients.

USE RESTRICTIONS

Do not apply NEGATE 37WG to turf under stress from drought, insects, disease, cold temperatures, high temperatures of above 85°F on cool season grasses, or poor fertility as injury may result.

Do not apply to residential lawns.

Do not apply to areas where children can contact treated turf.

Do not apply this product through any type of irrigation system.

Do not apply an organophosphate insecticide or nematicide within 7 days of a NEGATE 37WG application as injury potential to the desired grass may increase.

Do not apply to turf less than 1 year old.

Do not use on Bahiagrass where it is the desired turf, as severe injury may result.

Do not plant ornamentals such as shrubs and trees in treated areas for at least 1 year after the last application or bedding plants for at least 2 years.

Do not apply this product by air.

Do not use more than 4 applications of NEGATE 37WG per acre per year, or a maximum of 6 ounces product per acre per year. Please note the restrictions for the control of Bahiagrass,

DO NOT USE ON FOOD OR FEED CROPS.

Injury to trees or other desirable plants may result from failure to observe the following: Do not apply NEGATE 37WG (except as directed) or drain or flush equipment on or near trees or other desirable plants, or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

When overseeding, wait 2 months (8 weeks) after application. Do not apply to any body of water, including streams, irrigation water or wells. Do not apply where runoff water may flow onto agricultural land, as injury to crops may result. Do not allow spray drift onto adjacent crops or other desirable plants or trees as injury may occur.

See label for proper Mixing Instructions

Limitations, Restrictions, and Exceptions

Broadleaf (dicot) weeds

Dollarweed: A repeat application may be required in 4 to 6 weeks.

Canada thistle: Suppression only involving a visual reduction in competition compared to an untreated area.

Virginia buttonweed: Controls seedling Virginia buttonweed. Suppression only of more mature plants. Repeat application may be required in 4 to 6 weeks

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)