

TURF - DOLLAR (CURATIVE RATE)

General Information

GENERAL PRODUCT INFORMATION

Thiram Granuflo Agricultural Fungicide is a broad-spectrum fungicide for plant disease control in the culture of Peaches and Strawberries. Thiram Granuflo Agricultural Fungicide is a broad-spectrum fungicide for use on Turf (golf courses) to control Dollar Spot, Brown Patch and Fusarium Patch. When used as a Seed Protectant, Thiram Granuflo Agricultural Fungicide will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off or seedling blights. When used as a Repellent, Thiram Granuflo Agricultural Fungicide can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation.

Taminco Inc.'s recommendations for the use of this product are based upon tests believed to be reliable. **FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THE LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

Do not use on residential lawns or turf.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

For agricultural foliar applications use recommended rates for dilute, concentrate and aerial applications. Ground application is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo Agricultural Fungicide at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 20 gallons per acre. For aerial application, apply the recommended rate listed in

the \"per acre\" column of the Specific Applications table in a minimum of 10 gallons per acre. Applications of Thiram Granuflo Agricultural Fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage.

Make fresh spray mixtures for each application.

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers.

Limitations, Restrictions, and Exceptions

TURF APPLICATIONS:

Thiram Granuflo Agricultural Fungicide prevents infestation of common turf diseases such as Dollar Spot, Brown Patch and Fusarium Patch during the Summer growing season and Snow Mold during the cool weather seasons. When used at curative rates, Thiram Granuflo Agricultural Fungicide will cure diseased turf infected with Dollar Spot and Brown Patch and can also prevent Fusarium Patch when two applications are made during the Fall through the Spring season. Apply Thiram Granuflo Agricultural Fungicide with properly calibrated mechanical or motorized spray equipment such as ground boom.

DIRECTIONS FOR USE

Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. after disease symptoms first appear. Repeat at 7 to 10 day intervals as needed. Do not apply more than 40 oz per 1000 sq. ft. during the Summer growing season.

DO NOT graze or feed treated turf.

Golf Course Tees and Greens: The maximum winter application rate for turf is a single application of 16.3 lbs ai/acre or 0.374 lbs. ai/1000 sq ft. The maximum summer application rate for turf is 3 applications of 10.2 lbs. ai/acre or 0.234

ai/1000 sq ft. The maximum annual application rate is 47 lbs ai/acre or 1.08 lbs ai/1000 sq ft.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Exception: Once treated seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction, if there will be no contact with the treated seeds.

Timings

[After disease symptoms first appear.](#)