

FOR CONTROL OF CALIFORNIA GROUND SQUIRREL

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS:

Secure bait stations at locations where they will be readily accessible to target ground squirrels, preferably near active burrows. Secure stations 20 to 100 feet apart, depending upon the density of the California ground squirrel population. Use 2 to 4 pounds of bait per bait station.

Check bait stations frequently to insure that an uninterrupted supply of bait is always available. Maintain an uninterrupted supply of fresh bait for at least 30 days or until all signs of feeding have stopped. Collect and dispose of all dead animals and leftover bait properly.

Bait stations may be maintained in areas on permanent bases but should only be loaded with toxic bait when California ground squirrels are active and accepting grain-based foods in numbers sufficient to promote effective control.

USE RESTRICTIONS: This anticoagulant bait may be used to control California ground squirrels (*Spermophilus beecheyi*) in secured baitstation applications in or near ditch banks, along fence lines, and in noncrop areas. DO NOT broadcast this bait, apply it directly to food or feed crops, or use it in any manner other than that prescribed by the label. Wear protective gloves when applying bait. With detergent

and hot water, wash all implements used for applying bait. Do not use these implements for mixing, holding or transferring foods or feeds.

Method

[Directed](#)

[Bait](#)

Rates

[field_rates 0](#)

•

Timings

[N. A.](#)