

# **BLUEBERRIES (POST-HARVEST) - PHOMOPSIS CANE AND TWIG BLIGHT, MUMMYBERRY**

General Information

## **DIRECTIONS FOR USE**

SULFORIX is a balanced fungicide-insecticide-miticicide. For best results, spray to cover all exposed plant parts and repeat as necessary.

Not for residential use or application to residential sites. This product may not be used in, on, or around any structure, vehicle, article, surface or area associated with the household, including nonagricultural outbuildings, non-commercial greenhouses, pleasure boats and recreational vehicles; in or around any preschool or day care facility or on humans or pets.

Important Note: SULFORIX is a contact fungicide. Thorough coverage is necessary.

Do not use on apricots as they are susceptible to sulfur injury. SULFORIX may burn foliage when temperature is high (85°F). Do not make applications at such times. When applied to crops with foliage, let 30 days elapse between treatments with SULFORIX and treatments with an oil spray. Do not combine SULFORIX with other pesticides unless previous experience has shown them to be compatible. If tank mixing, always follow the more restrictive label directions. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Aerial applications of SULFORIX are prohibited.

If crust occurs on surface of SULFORIX, break through. Crystallization does not impair product quality.

Do not apply, or allow to drift, to painted surfaces as a permanent stain may result.

Do not apply this product through any type of irrigation system.

Do not apply more than 30 gallons of this product/acre/year to any crop listed on the label.

## Limitations, Restrictions, and Exceptions

### BLUEBERRIES

Apply POST HARVEST (September or October) and/or as a DELAYED DORMANT spray of 1 to 2 gallons of SULFORIX per 100 to 150 gallons of diluted spray per acre. During the growing season, apply SULFORIX at 2 quarts per acre in a minimum of 50 gallons of finished spray per acre. Do not make multiple applications within 10 days of each other.

NOTE TO USER: Do not use within 14 days of an oil spray or when temperatures are above 85°F. Burning of foliage may occur during periods of warm temperatures.

### Method

[Broadcast/Foliar Ground](#)

### Rates

[field\\_rates 0](#)

- 

### Restricted Entry Interval

48 hours

### Timings

[Post-harvest](#)