

TREE NUTS - APHIDS

General Information

USE INFORMATION

APTA insecticide is a suspension concentrate containing 1.31 lbs of active ingredient tolfenpyrad per gallon. This product is a contact insecticide used for the control of several orders of insects. Complete and thorough spray coverage is necessary for maximum results. APTA insecticide should be used in a program with other products to provide season long protection. Apply as a spray as directed in the Application Directions section of the label.

Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. Adjust water volumes and tractor speed accordingly for crops with dense foliage or excessive growth. Unless otherwise specified under Application Directions, apply when pest populations are beginning to build, before crop damage or injury is observed. Consult your local agricultural advisor or state cooperative extension service for recommendations.

DIRECTIONS FOR USE OF APTA INSECTICIDE AS A FUNGICIDE

For crops and diseases where the level of activity of APTA insecticide is listed as “control”, this product may be used alone as a contact fungicide or mixed with other registered fungicide products to broaden spectrum of disease control. For crops and diseases where the level of activity of APTA insecticide is listed as “suppression”, this product should NOT be substituted for labeled fungicidal products.

APPLICATION DIRECTIONS

- Applications should be made immediately after the spray solution is prepared.
- Thorough spray coverage is critical to obtain control of the target pest(s).
- Applications may be made by ground with high or low volume spray equipment that provides thorough spray coverage of the plant.

- For ground applications, use coarse droplet size.
- Use sufficient water volume to ensure thorough coverage of foliage.
- Do not apply APTA insecticide through any type of irrigation system.
- RESTRICTION: Not for sale or use in the state of New York.

Limitations, Restrictions, and Exceptions

USE RESTRICTIONS

- Do not apply by air.
- Maintain a minimum of 25 feet of vegetative filter (buffer) strip.
- For ground applications, do not use less than 50 gallons of water per acre.
- Do not apply by Alternate Row Middle (ARM) spray method.
- Allow at least 14 days between applications.

USE RECOMMENDATIONS

- Use sufficient water volume to ensure thorough coverage of foliage. Thorough spray coverage is critical to obtain control of the target pest(s).

Method

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[When pest populations are beginning to build.](#)