

CORN, SWEET

General Information

General Information

Stamina F3 HL is a liquid seed treatment used for control or suppression of certain seedborne and soilborne diseases of wheat and barley.

Stamina F3 HL will generally increase emergence and plant stands by controlling or suppressing these diseases.

Method of Application

Stamina F3 HL is for use at commercial seed treatment facilities using standard commercial seed treatment equipment (such as, but not limited to, slurry or mist-type equipment). Not for use on farm. Seed should be well conditioned and free of chaff before treating with Stamina F3 HL. Treated seed should not require drying after treatment and can be stored or bagged immediately (provided excessive slurry volumes are not applied to the seed). Stamina F3 HL should be applied to wheat and barley at the rate of 1.0 fluid ounce of product per 100 pounds of seed. Thorough seed coverage and proper application equipment calibration is essential for good disease control. To achieve thorough coverage, Stamina F3 HL may be diluted with water, if needed, into a slurry. Direct application into a planter box is prohibited. Use an EPA-approved dye that imparts an unnatural color to the seed. Any dye or colorant added to treated seed must be cleared for use under 40 CFR Section 180.900.

Use Restrictions

- DO NOT use treated seed for food, feed or oil processing.
- DO NOT store Stamina F3 HL near feed or foodstuffs.
- DO NOT contaminate feed or foodstuffs.
- DO NOT contaminate domestic or irrigation waters.
- DO NOT plant any crop not listed on the label within 30 days after planting

Stamina F3 HL fungicide treated seed.

- DO NOT use the same auger for treating seed that will handle commercial or feed grains.

- DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed. Store treated seed in cool, dry conditions.

Product Information

Stamina F3 HL is a liquid seed treatment used for control or suppression of certain seedborne and soilborne diseases of corn, oats, rye, sorghum and triticale. Stamina F3 HL may increase emergence and plant stands by controlling or suppressing these diseases.

Seed treatment applications with Stamina F3 HL have been associated with more rapid and increased emergence of seedlings under certain cold conditions. Make all applications according to the use directions in the Stamina F3 HL main label. Use the higher rates of Stamina F3 HL when disease pressure is expected to be high.

Limitations, Restrictions, and Exceptions

Rate

Equivalent to 0.0338 mg ai/kernel and 0.44 fl oz product per 80,000 kernel count (based on 1818 kernels per pound)

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

-

Timings

[N.A.](#)