

PECANS - PECAN NUT CASEBEARER

General Information

GENERAL INFORMATION

Confirm 2F insecticide mimics the action of the natural insect hormone 20-hydroxyecdysone, the physiological inducer of the molting and metamorphosis process in insects. Confirm 2F is highly active against most lepidopterous larvae while having practically no activity at typical use rates against other orders of insects. The selectivity of Confirm 2F allows for the maintenance of the populations of beneficial and predatory insects, which is a key element in integrated pest management programs. Confirm 2F controls lepidopterous larvae through a novel mode of action by the induction of a premature lethal molt which initiates within hours of ingestion of treated crop surfaces. Contact activity has also been observed in some insects. Actual death of the larvae will take several days to occur, although feeding by the insects generally ceases within 24 hours of ingestion.

Application Timing

The activity of Confirm 2F is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent upon the feeding behavior of the target pest. For internal feeding larvae, application must be made prior to the time that surface feeding occurs. For foliar or surface feeding larvae, application made while active feeding is occurring will be effective.

Reapplication may be required to protect new flushes of foliage or rapidly expanding fruit. The reapplication interval will vary depending upon how rapidly the crop is growing and the generation time of the target pest. While Confirm 2F is essentially equally effective against all instars, it is generally good practice to make applications to early instars to avoid the heavy damage that can be inflicted by later instar larvae. For best results, begin applications when first signs of feeding damage or when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities, to determine the appropriate threshold for application in your area.

Limitations, Restrictions, and Exceptions

APPLICATION TIMING

For each generation, apply at the initiation of egg hatch (for first generation this is approximately 8 to 15 days following first sustained moth catch¹. Control of first generation pecan nut casebearer may require a second application under conditions of extended egg lay or for improved coverage of rapidly expanding nuts and foliage. Use higher rates for extended residual effectiveness, higher pest infestations, low crop load, larger trees or heavy, dense foliage.

RESTRICTIONS

- Do not apply more than 122 fl oz per season.
- Allow at least 14 days to elapse between final application and harvest.

Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

Ground Application: Make applications of Confirm 2F by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre.

Aerial Application: Make application of Confirm 2F in a minimum of 5 gallons per acre. Confirm 2F can be applied by aerial applications when situations warrant. However, application by this method may result in reduced efficacy if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

First sustained moth catch (biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period. Consult State Extension Specialists or other qualified authorities for specific information regarding number, placement and management of pheromone traps.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

4 hours

Timings

[At the initiation of egg hatch.](#)