

# **CEREAL GRAINS - BARLEY, BUCKWHEAT, ETC.**

## General Information

### PRODUCT INFORMATION

Dyna-Shield Fludioxonil is a fungicide for seed treatment use to control soil-borne and seed-borne organisms such as *Aspergillus*, *Fusarium*, *Helminthosporium*, *Penicillium*, and *Rhizoctonia* that cause seed decay, damping off, and seedling blights.

If a rate range is indicated in the product instructions below, use the higher rate when high disease pressure is anticipated based on conditions and experience for your area.

### USE RESTRICTIONS

- Store treated seed away from food and feedstuffs.
- DO NOT allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- DO NOT contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed package or containers in accordance with local requirements.
- For seeds that are treated and bagged for later use the following must appear on the bag:
  - This seed has been treated with fludioxonil seed treatment.

- DO NOT use treated seed for feed, food or oil purposes.

### Seed Treatment (Export Use)

FOR TREATMENT OF THE SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH DYNA-SHIELD FLUDIOXONIL AND DYNA-SHIELD METALAXYL BEFORE IMPORTATION.

User should verify actual registered rate in country of destination and/or use. Seed treated in accordance with the export use pattern must be labeled as follows:

TREATED SEED: FOR EXPORT ONLY - NOT FOR DOMESTIC SALE OR USE. TREATED WITH THE FUNGICIDE, FLUDIOXONIL

Dyna-Shield Fludioxonil - Use Dyna-Shield Metalaxyl or an equivalent rate of an appropriate metalaxyl or mefenoxam formulation.

### Limitations, Restrictions, and Exceptions

Barley, buckwheat, millet (pearl and Proso), oats, rye, teosinte, triticale, wheat, and wild rice

Target Disease: Decay, damping-off, and seedling blight caused by seed-borne and soil-borne fungi.

- Dyna-Shield Fludioxonil contains 14.18 grams active ingredient per fluid ounce.
- Cereal forage may not be grazed until 30 days after planting.

### Method

[Seed Treatment](#)

Rates

[field rates 0](#)

- 

### Restricted Entry Interval

12 hours

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact

with anything that has been treated.

Timings

[N.A.](#)