

CEREAL GRAINS - CORN (FIELD, POP)

General Information

PRODUCT INFORMATION

Dyna-Shield Fludioxonil is a fungicide for seed treatment use to control soil-borne and seed-borne organisms such as *Aspergillus*, *Fusarium*, *Helminthosporium*, *Penicillium*, and *Rhizoctonia* that cause seed decay, damping off, and seedling blights.

If a rate range is indicated in the product instructions below, use the higher rate when high disease pressure is anticipated based on conditions and experience for your area.

USE RESTRICTIONS

- Store treated seed away from food and feedstuffs.
- DO NOT allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- DO NOT contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed package or containers in accordance with local requirements.
- For seeds that are treated and bagged for later use the following must appear on the bag:
 - This seed has been treated with fludioxonil seed treatment.

- DO NOT use treated seed for feed, food or oil purposes.

Seed Treatment (Export Use)

FOR TREATMENT OF THE SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH DYNA-SHIELD FLUDIOXONIL AND DYNA-SHIELD METALAXYL BEFORE IMPORTATION.

User should verify actual registered rate in country of destination and/or use. Seed treated in accordance with the export use pattern must be labeled as follows:

TREATED SEED: FOR EXPORT ONLY - NOT FOR DOMESTIC SALE OR USE. TREATED WITH THE FUNGICIDE, FLUDIOXONIL

Dyna-Shield Fludioxonil - Use Dyna-Shield Metalaxyl or an equivalent rate of an appropriate metalaxyl or mefenoxam formulation.

Limitations, Restrictions, and Exceptions

Corn (field, pop)

For Pythium spp. control, use Dyna-Shield Fludioxonil in combination with labeled rates of Dyna-Shield Metalaxyl. Observe all precautions, limitations, rates, and directions for use on the respective labels before applying.

Dyna-Shield Fludioxonil will also control seed-borne head smut caused by Sphacelotheca reiliana.

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N.A.](#)