

# **GRASS FORAGE, FODDER AND HAY**

## General Information

### PRODUCT INFORMATION

Dyna-Shield Fludioxonil is a fungicide for seed treatment use to control soil-borne and seed-borne organisms such as *Aspergillus*, *Fusarium*, *Helminthosporium*, *Penicillium*, and *Rhizoctonia* that cause seed decay, damping off, and seedling blights.

If a rate range is indicated in the product instructions below, use the higher rate when high disease pressure is anticipated based on conditions and experience for your area.

### USE RESTRICTIONS

- Store treated seed away from food and feedstuffs.
- DO NOT allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- DO NOT contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed package or containers in accordance with local requirements.
- For seeds that are treated and bagged for later use the following must appear on the bag:
  - This seed has been treated with fludioxonil seed treatment.

- DO NOT use treated seed for feed, food or oil purposes.

### Seed Treatment (Export Use)

FOR TREATMENT OF THE SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH DYNA-SHIELD FLUDIOXONIL AND DYNA-SHIELD METALAXYL BEFORE IMPORTATION.

User should verify actual registered rate in country of destination and/or use. Seed treated in accordance with the export use pattern must be labeled as follows:

TREATED SEED: FOR EXPORT ONLY - NOT FOR DOMESTIC SALE OR USE. TREATED WITH THE FUNGICIDE, FLUDIOXONIL

Dyna-Shield Fludioxonil - Use Dyna-Shield Metalaxyl or an equivalent rate of an appropriate metalaxyl or mefenoxam formulation.

### Limitations, Restrictions, and Exceptions

Grass Forage, Fodder and Hay: all grasses (including turfgrass, bermudagrass, bluegrass, and fescue or bromegrass), Graminae family (either green or cured) except sugarcane and those included in the cereal grains group that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage.

Note: Forage may not be grazed until 30 days after planting.

### Use Rate

(Fl Oz of Product/100 Lb of Seed, Propagating Root, Tuber Material)

### Method

[Seed Treatment](#)

### Rates

[field rates 0](#)

- 

### Restricted Entry Interval

12 hours

Exception: If the seed is treated with the product and the treated seed is soil-

injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N.A.](#)