

# **SPOT TREATMENT OF COMMERCIAL SOD FARMS**

## General Information

### GENERAL INFORMATION

Foliage sprays should be applied during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. Use low spray pressures to minimize spray drift. Apply this herbicide in a manner to avoid contacting nearby susceptible crops or other desirable plants and to avoid contaminating water intended for irrigation or domestic use. Read and follow all use precautions given on the label.

### GENERAL PRECAUTIONS and RESTRICTIONS

In Arizona: The state of Arizona has not approved Chaser Turf Herbicide for use on sod farms.

For ground application only.

### Weeds Controlled

Bull thistle, Burdock, Buttercup, Canada thistle, Mustards, Wild carrot, Cornspeedwell, Ground ivy, Oxalis (stricta and corniculata), Prostrate spurge and Wild violet: See Broadcast Treatment of Residential, Commercial, and Recreational Turf and Commercial Sod Farms\".

### Limitations, Restrictions, and Exceptions

#### Spot Treatment of Residential, Commercial, and Recreational Turf

Mix 1.0 to 2.0 fluid ounces (0.01 to 0.03 pound acid equivalent) of Chaser Turf Herbicide in enough water to make 3.0 gallons of spray (1.0 to 2.0 quarts per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of undesirable plants to point of runoff.

### Method

[Spot treatment](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

[field\\_rates 2](#)

- 

Restricted Entry Interval

12 hours

Timings

[At any time broadleaf weeds are susceptible.](#)