

# WALNUTS

## General Information

### DIRECTIONS FOR USE

Avoid application of oil sprays just prior to or during hot weather, rainy weather or drying winds. Apply when temperature is above 40°F. Do not apply just prior to freezing weather. Use only when trees are in healthy condition, and when moisture conditions are sufficient. Avoid overspraying.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Do not apply this product through any type of irrigation system.

Do not use if material does not emulsify. Do not use in undiluted form.

Apply the following specified rates in sufficient water to thoroughly cover one acre. With ground application equipment on Apples and Pears use a minimum of 300 gallons per acre. On stone fruits and nuts use a minimum of 200 gallons per acre. On grapes for cane pruned vines use a minimum of 200 gallons per acre. For cordon pruned or arbor trained vines use a minimum of 300 gallons of diluted spray per acre. When using concentrate spray equipment use a minimum of 60 to 90 gallons of diluted spray per acre.

### Limitations, Restrictions, and Exceptions

#### WALNUTS

Use 3 gallons per 100 gallons of water. Apply in the delayed dormant period. This treatment should only be used on trees that have not suffered from lack of moisture at anytime during the growing season. Further, oil should not be applied to orchards when soil moisture is low.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

4 hours

Timings

[Delayed dormant period.](#)