

RICE - APPLICATION FOR SOUTHERN UNITED STATES - GRASS WEEDS

General Information

PRODUCT INFORMATION

Forfeit 280 is a water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds in LibertyLink canola, LibertyLink corn, LibertyLink cotton, and LibertyLink soybean, and in trees, vines, and berries. Forfeit 280 may be applied for potato vine desiccation. Forfeit 280 may also be applied as a broadcast burndown application before planting or prior to emergence of any conventional or transgenic variety of canola, sweet corn*, corn, cotton, olive, rice*, soybean, or sugar beet.

*Not for this use in California.

Forfeit 280 is only foliar-active with little or no activity in soil. Weeds that emerge after application will not be controlled. Apply Forfeit 280 to actively growing weeds as described in the Weed Control Recommendations for Row Crops section to get maximum weed control.

Uniform, thorough spray coverage is necessary to achieve consistent weed control. Necrosis of leaves and young shoots occur within 2 to 4 days after application under good growing conditions.

- Forfeit 280 is rainfast 4 hours after application to most weed species; therefore, rainfall within 4 hours may necessitate retreatment or may result in reduced weed control.

- Applications should be made between dawn and 2 hours before sunset to avoid the possibility of reduced lambsquarters and velvetleaf control.

- Consult your local Cooperative Extension Service or Loveland Products, Inc. representative for guidelines on the optimum application timing for Forfeit 280 in your region.

- Weed control may be reduced if application is made when heavy dew, fog, and

mist/rain are present; or when weeds are under stress due to environmental conditions such as drought, cool temperatures, or extended periods of cloudiness.

- To maximize weed control, do not cultivate from 5 days before an application to 7 days after an application.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS FOR USE ON RICE

Application Timing for the Southern United States (Arkansas, Louisiana, Mississippi, Missouri, Texas)

Applications of Forfeit 280 on rice may be made from the 1-leaf stage through the mid-tillering stage of development. Refer to the Rate Tables for Weed Control in Rice to select the proper rate to use to control the weed species present. Forfeit 280 will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.

Rice fields should be as level as possible and free of large clods to obtain uniform germination of rice and grassy weeds and to ensure uniform flood levels. If necessary, fields may be flushed prior to treatment so that the rice and grass/broad leaf weeds are actively growing at the time of treatment. If the rice field is flushed, allow sufficient time for germination of the weed species to occur prior to treatment.

Apply Forfeit 280 prior to the permanent flood when weeds are in the 1- to 5-leaf stage. A second application is recommended after a new flush of weeds emerge. A second application may be made from 10 days after the first application up to the mid-tillering growth stage of the rice. For optimum weed control, apply Forfeit 280 before canopy closure to ensure thorough spray coverage of the weed species.

When applying Forfeit 280 post-flood, lower the water level so that 75% of the weed foliage is exposed. The water level may be brought back to normal level 48 hours after the herbicide application.

NOTE:

For optimum red rice control, make 2 applications of Forfeit 280. The first application should be made when the red rice is in the 2- to 3-leaf stage. The second application should be made after the newly emerged red rice reaches the 2-

to 3-leaf stage, but before the white rice reaches the mid-tillering stage of development.

Restrictions to the Directions For Use on Rice

- DO NOT plant rotation crops in a field treated with Forfeit 280 within 120 days after the last application of Forfeit 280 with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of Forfeit 280. The crops listed on the label may be planted at any time.
- DO NOT apply Forfeit 280 through any type of irrigation system.
- DO NOT use paddy water from a rice field treated with Forfeit 280 for irrigation, or as a water source for livestock or for raising crayfish.
- DO NOT add surfactants or crop oils. A silicon-based anti-foam agent may be added if needed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

70 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Exception of sweet corn irrigation activities, which has a 4-day REI.

Timings

[Postemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)