

TURF, ORNAMENTAL - BLACK MEDIC, CARPETWEED, ETC.

General Information

PRODUCT INFORMATION

Chaser 2 Amine herbicide is for use on Fence Rows, Non-Irrigation ditchbanks, Roadsides, other Non-crop Areas, Industrial sites, Rights-of-way.

USE PRECAUTIONS

Be sure that use of this product conforms to all applicable regulations.

Restrictions on Retreatment: Do not apply more than 2 broadcast applications per year per treatment site.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not apply Chaser 2 Amine directly to, or otherwise permit it to come into direct contact with, cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants and do not permit spray mists containing it to drift onto them. Very small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

Do not use for manufacturing or formulating.

Do not apply to exposed roots of shallow rooted trees or shrubs.

Do not apply on ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

USE PRECAUTIONS

Foliar sprays should be applied during warm weather when brush and weeds are actively growing. Application under drought conditions may provide less than desirable results.

With aerial applications use a drift control system such as Microfoil or Thru-Valve

booms, or high quality drift control additive. Keep spray pressures low enough to provide coarse spray droplets. Spray booms should be no longer than $\frac{3}{4}$ of the rotor length. Do not use a thickening agent with the Microfoil or the Thru-Valve booms, or other systems that cannot accommodate thick sprays. Spray only when the wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to air inversions.

NOTE: Do not use spray equipment for other applications to land planted, or to be planted to susceptible crops or desirable plants, unless it has been determined that all phytotoxic herbicide residue has been removed by thorough cleaning of the equipment.

APPROVED USES

Grazing Restrictions

If site is used for grazing:

Apply a maximum of 1 lb per acre Triclopyr (1 gal. of Chaser 2 Amine) per application.

Do not make more than one application per year.

Do not graze lactating dairy cattle until the next growing season after application.

Remove meat animals from treated areas 3 days prior to slaughter.

Limitations, Restrictions, and Exceptions

Turf, ornamental (golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas)

Postemergence: Limited to 2 applications per year. Maximum of 1.5 lbs ae 2,4-D per acre per application. The maximum seasonal rate is 3.0 lbs ae 2,4-D per acre, excluding spot treatments.

Method

[Spray](#)

Rates

[field rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

-

Restricted Entry Interval

48 hours

Timings

[Postemergence \(Weed\)](#)