

SOYBEANS

General Information

PRODUCT INFORMATION

ENHANCE Seed Protectant is a formulation specifically designed for treatment of wheat, oat and barley seed on the farm at planting time. It controls covered smut (*Tilletia caries* and *Tilletia foetida*) and loose smut (*Ustilago nuda*) on wheat, covered smut (*Ustilago kollerii*) and loose smut (*Ustilago avenae*) on oats and covered smut (*Ustilago hordei*) and loose smut (*Ustilago nuda*) on barley. It also tends to reduce seed rot and seedling blight caused by soil fungi.

ENHANCE also protects soybean seed from seed borne and soil borne fungi that cause seed decay, damping off, and seedling blights (including *Fusarium*, *Rhizoctonia* and *Pythium*). For use only on agricultural establishments at or immediately before planting.

RESTRICTIONS

- Do not use treated seed for food, feed, or oil purposes.
- Do not graze or feed livestock on treated area for 42 days after planting.
- Soybean restriction: Do not graze or feed forage or hay from treated areas to livestock.

APPLICATION INSTRUCTIONS

Apply as a planter-box treatment (including air and vacuum planters), mixing thoroughly with the seed before planting. For best results, the following procedure is suggested. Fill planter-box only half-full of seed, add half the required amount of ENHANCE and mix thoroughly with a stick. Then add remainder of seed to planter-box and add the rest of the required amount of product to the seed, again mixing thoroughly. Good results are also obtained by premixing seed with the product in a suitable container, then pouring treated seed into planter-box.

Note: One level scoop (enclosed) is equivalent to approximately 1 avoirdupois ounce of this product.

Adjust the seed drill or planter to give the required rate of seed delivery with treated seed. Regularly inspect planter plates and clean as required. At the end of

each day, clean out the electric eye monitors as directed by the manufacturer.

Limitations, Restrictions, and Exceptions

RESTRICTIONS

- Do not use treated seed for food, feed, or oil purposes.
- Do not graze or feed livestock on treated area for 42 days after planting.
- Do not graze or feed forage or hay from treated areas to livestock.

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard, allows workers to enter the treated areas without restriction if there will be no contact with the soil/media subsurface.

Timings

[Preplant](#)