

ORNAMENTALS

General Information

GENERAL INFORMATION

HI-YI ELD GRU B FREE ZONE will control insects on turf and ornamentals grown in residential landscapes. HI-YIELD GRUB FREE ZONE is a systemic insecticide applied as a soil treatment only and is translocated upward within the plant. For best results, HI-YIELD GRUB FREE ZONE must be applied where the roots of the target plants can absorb the active ingredient. After application, irrigate moderately but thoroughly being sure to avoid leaching and runoff from the container for at least three irrigations or 10 days whichever is longer.

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

HI-YIELD GRUB FREE ZONE General Use Precautions:

- Do NOT apply more than 1.8 lbs (0.009 lb. of active ingredient) per 1,000 sq. ft. per year.
- Following application, avoid runoff or puddling of irrigation water.
- Avoid application of HI-YIELD GRUB FREE ZONE to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant.

Limitations, Restrictions, and Exceptions

Application to Ornamentals

GRUB FREE ZONE may NOT be used in commercial greenhouses or nurseries.

HI-YIELD GRUB FREE ZONE can be applied as a broadcast application on ornamentals intended for aesthetic purposes or climatic modification in residential landscapes such as interior plantscapes, ornamental gardens or lawns and grounds.

The need for an application can be used on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made preceding or during the egg laying period.

Specific Instructions

Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.

For optimum control, mechanical incorporation, irrigation or rainfall should occur within 24 hours after application to move the active ingredient into soil.

Be sure to read the “Application Equipment” section of the label under “GENERAL INFORMATION” above.

Method

[Broadcast](#)

Rates

[field rates 0](#)

-

Timings

[Preceding or during the egg laying period.](#)