

ROSE - BLACK SPOT, POWDERY MILDEW, RUST

General Information

Do not allow people or pets to contact treated plants until spray has dried.

Do not water the treated area to the point of runoff.

Do not make applications during rain.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures
2. Applications to lawns, turf and other vegetation All outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

USE TIPS

- When using this product on plants not listed, test on a small area first.
- Do not spray if temperature is above 90°F as damage to some delicate plants can occur.
- Spray early in the morning or late in the evening, when air is still.
- If it rains before product has dried, reapply.
- Regular light sprayings are more effective than a single heavy treatment.

SENSITIVE PLANT CARE

Use with care on Ferns, Palms and Fuchsia. As with all plants, especially Carnations,

Chrysanthemums and Roses, there exists a wide variety of species and, therefore, a wide variety of susceptibility to injury. To test for plant sensitivity, spray a few plants as described above, wait 24 hours and check for possible injury.

Limitations, Restrictions, and Exceptions

Rose: Treat on a seven- to 10-day schedule. If Black spot is not a problem, a 10- to 14-day schedule is recommended.

Method

[Spot treatment](#)

Timings

[Early in the morning or late in the evening.](#)