

BANANA, PLANTAIN

General Information

PRODUCT INFORMATION

Spray Volume: Apply REPAR TEBUCON 45 DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under “REMARKS” for each crop. Increase spray volume as crop growth increases to ensure through coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

Limitations, Restrictions, and Exceptions

BANANA, PLANTAIN

Apply specified dosage as a foliar spray in a minimum of 20 gallons or more of water per acre to assure thorough coverage of the plants. A spray oil recommended for use on banana or plantain should be utilized at 4.3 pints per acre in tank-mix with REPAR TEBUCON 45 DF FUNGICIDE. Thoroughly mix REPAR TEBUCON 45 DF FUNGICIDE in the spray solution prior to the addition of a spray oil. Make the first application before disease becomes established and repeat applications at 14 -day intervals if necessary to maintain disease control.

Do not apply REPAR TEBUCON 45 DF FUNGICIDE to non-bagged bananas. A maximum of 16 ounces of REPAR TEBUCON 45 DF FUNGICIDE maybe applied per acre per harvest cycle. REPAR TEBUCON 45 DF FUNGICIDE may be applied up to the day of harvest.

Method

[Foliar spray](#)

Pre-Harvest Interval

14 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Before disease becomes established.](#)