

PISTACHIO

General Information

PRODUCT INFORMATION

Spray Volume: Apply REPAR TEBUCON 45 DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under “REMARKS” for each crop. Increase spray volume as crop growth increases to ensure through coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

Limitations, Restrictions, and Exceptions

PISTACHIO

Botryosphaeria: Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of REPAR TEBUCON 45 DF FUNGICIDE 14 days apart beginning 49 days before harvest.

APPLICATION RESTRICTIONS: Apply REPAR TEBUCON 45 DF FUNGICIDE in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.

A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. REPAR TEBUCON 45 DF FUNGICIDE may be applied up to 35 days before harvest.

Control of insect vectors and a thorough pruning program to remove plant tissue

infected by Botryosphaeria are critical for optimum control of this disease.

RESISTANCE MANAGEMENT STRATEGY: REPAR TEBUCON 45 DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing REPAR TEBUCON 45 DF FUNGICIDE with other DMI fungicides is not recommended.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

35 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[At 25 to 50% bloom.](#)

[To protect young flower clusters and fruit.](#)