

## **FILBERT (HAZELNUT)**

### General Information

#### PRODUCT INFORMATION

Spray Volume: Apply REPAR TEBUCON 45 DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under “REMARKS” for each crop. Increase spray volume as crop growth increases to ensure thorough coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

#### Limitations, Restrictions, and Exceptions

#### FILBERT

Apply REPAR TEBUCON 45 DF FUNGICIDE in a preventive spray schedule when conditions are favorable for disease (such as budbreak when the first green tissue is visible). Continue applications of REPAR TEBUCON 45 DF FUNGICIDE at 7- to 14-day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during budbreak and rapid shoot elongation. The expanding shoot tip must be protected.

Reduce the application interval for varieties that are highly susceptible to the indicated disease or when severe disease conditions exist.

Apply REPAR TEBUCON 45 DF FUNGICIDE in sufficient spray volume for thorough coverage. Tankmixing the lowest labeled rate of a spray surfactant with REPAR TEBUCON 45 DF FUNGICIDE may improve spray coverage and penetration of the active ingredient into plant tissue. The use of ground application is preferred because of the difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and stems by air. A maximum of 32 ounces of REPAR TEBUCON 45 DF FUNGICIDE may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. REPAR TEBUCON

45 DF FUNGICIDE may be applied up to 35 days before harvest.

REPAR TEBUCON 45 DF FUNGICIDE is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing REPAR TEBUCON 45 DF FUNGICIDE with other DMI fungicides is not recommended.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

35 days

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[When conditions are favorable for disease \(such as budbreak when the first green tissue is visible\).](#)