

POME FRUITS

General Information

PRODUCT INFORMATION

Spray Volume: Apply REPAR TEBUCON 45 DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under “REMARKS” for each crop. Increase spray volume as crop growth increases to ensure through coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

Limitations, Restrictions, and Exceptions

POME FRUITS

Scab:

Preventive Schedule: Apply at 7- to 10-day intervals from green tip through the second cover spray.

Post-Infection Schedule: Apply within 72 hours after the beginning of a scab infection period. REPAR TEBUCON 45 DF FUNGICIDE applications should not be closer than 7 days apart. Continue REPAR TEBUCON 45 DF FUNGICIDE applications using the preventive schedule if the post-infection schedule is discontinued.

Cedar-apple rust: Apply at 7- to 10-day intervals from pink stage through the second cover spray.

Powdery mildew: Apply at 7- to 10-day intervals from tight cluster through the second cover spray.

APPLICATION RESTRICTIONS:

- The amount of REPAR TEBUCON 45 DF FUNGICIDE required per acre will depend

on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated.

- For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

REPAR TEBUCON 45 DF FUNGICIDE may be applied up to 75 days before harvest. A total of 3 pounds of REPAR TEBUCON 45 DF FUNGICIDE may be applied per acre per crop season.

RESISTANCE MANAGEMENT STRATEGY: REPAR TEBUCON 45 DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Do not mix or alternate REPAR TEBUCON 45 DF FUNGICIDE with other DMI fungicides.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

75 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

5 days

Timings

[When conditions are favorable for disease \(such as budbreak when the first green tissue is visible\).](#)