

# PLUM

## General Information

### PRODUCT INFORMATION

Spray Volume: Apply REPAR TEBUCON 45 DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under “REMARKS” for each crop. Increase spray volume as crop growth increases to ensure through coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

### Limitations, Restrictions, and Exceptions

#### PLUM

Mix specified rate of REPAR TEBUCON 45 DF FUNGICIDE in 100 gallons of water, with or without wax. Maintain REPAR TEBUCON 45 DF FUNGICIDE suspension by direct or by-pass agitation sufficient to prevent settling of product in spray mixture. Apply as spray to fruit as they travel along packing line. After sizing, sorting or washing, REPAR TEBUCON 45 DF FUNGICIDE application may be incorporated into a waxing system using conventional equipment that provides for even and complete coverage of fruit. Do not apply more than once prior to boxing and do not rise.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field rates 0](#)

- 

#### Timings

[Post-harvest](#)