

TURFGRASS

General Information

DIRECTIONS FOR USE

When applying multiple products containing Atrazine active ingredient (a.i.), the total pounds of atrazine a.i. applied in pounds of atrazine a.i. per acre may not exceed 4 pounds of atrazine a.i. from all applications made in a year.

Turfgrass at Residential Sites (including homes, daycare facilities, schools, playgrounds, parks, recreational areas, and sports fields.) This product must be watered in immediately after application. Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing:

- This product must be watered in immediately.
- Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.atrazine-watershed.info or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Turf Care Supply Corp. for a refund.

IMPORTANT: In addition to providing turfgrass with required plant nutrients this product will control both \"weeds that are present\" and \"weeds germinating from seeds\".

FOR WEEDS THAT ARE PRESENT: Includes broadleaf weeds from the list such as:

Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.

FOR WEEDS GERMINATING FROM SEEDS: Applied before weed seeds germinate, includes grassy weeds from the list such as: Crabgrass, Paspalum, and Spotted Spurge.

WHERE TO USE

FOR CONTROL of weeds in St. Augustine and Centipede grass in or on residential lawns, corporate lawns, parks, recreational areas, golf course, cemeteries, and athletic fields.

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

NOT FOR USE on Bluegrass, Fescue, Bentgrass, Bahiagrass, Ryegrass, Bermudagrass, or Dichondra lawns.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post-emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Apply in early Spring for best control (January 1st - March 1st), otherwise it may be applied up to June 15th with good weed control.

For pre-emergence weed control apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

USE PRECAUTIONS

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

DO NOT APPLY THIS PRODUCT IN A WAY THAT WILL CONTACT ANY PERSON OR PET, EITHER DIRECTLY OR THROUGH DRIFT. KEEP PEOPLE AND PETS OUT OF THE AREA DURING APPLICATION. THIS PRODUCT MUST BE WATERED IN IMMEDIATELY AFTER APPLICATION. DO NOT ENTER OR ALLOW OTHERS (INCLUDING CHILDREN OR PETS) TO ENTER THE TREATED AREAS (EXCEPT THOSE INVOLVED IN THE WATERING) UNTIL THE WATERING-IN IS COMPLETE AND THE SURFACE IS DRY.

Limitations, Restrictions, and Exceptions

TURFGRASS

Apply 3.74 lbs to 5.00 lbs product uniformly over 1,000 sq ft of lawn area (equivalent to 1.5 - 2.0 lb Atrazine a.i. per acre). Use the lower rate for average weed problems; the higher rate for heavy weed populations. Do not exceed listed rates. Do not apply more than 4 lb a.i. Atrazine (2 applications at 5.00 lbs product rate) per acre per year. Apply to dry grass. This product must be watered in immediately after application.

Application with rotary or pendulum spreaders is suggested for uniform distribution. Consult your spreader manual or spreader manufacturer to determine proper application rates and spreader settings. You may need to adjust the setting depending on the walking speed, spreader condition, and product.

After application, sweep or blow any residual fertilizer on paved or other surfaces back onto the turf area.

Method

[N. A.](#)

Rates

[field_rates 0](#)

field_rates 1

-

Timings

Preemergence (Weed)

Postemergence (Weed)