

ORNAMENTAL TREES AND SHRUBS - CONIFER

General Information

HOW TO MIX AND APPLY

Unless otherwise directed, mix the recommended amount of HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS per 4 gallons of water and apply in accordance with specific Directions For Use.

Where appropriate, calculate the area to be treated by measuring the length and the width then multiply together to determine the square footage of the area to be treated.

Unless otherwise directed, apply HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS with a pump-up or other commercial type sprayer capable of delivering a coarse spray. When mixing, first fill the sprayer with half of the required amount of water, then measure and add the appropriate amount of HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS. Complete filling the sprayer with the remaining amount of required water. Shake the sprayer or initiate agitation to mix the product and water. Thorough spray coverage is necessary, but avoid spray runoff.

For large area coverage, this product may also be used through a hose-end type sprayer such as a ferti lome Air-O-Matic Hose-end Spray. Always follow the directions provided with the hose-end sprayer type for proper dilution and application of HIYIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS.

Do not use food utensils, such as measuring spoons or cups, for food purposes after use in measuring pesticides.

Limitations, Restrictions, and Exceptions

ORNAMENTAL TREES & SHRUBS

(IN RESIDENTIAL LANDSCAPED AREAS AND LANDSCAPED AREAS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS AND ATHLETIC FIELDS (INCLUDING TREES, SHRUBS, FLOWERS, EVERGREENS, FOLIAGE PLANTS AND GROUND COVERS)):

Not for use in Nurseries. Apply as a uniform spray, thoroughly wetting leaves and branches. Do not spray to the point of runoff. Apply to cover underside of leaves and to penetrate dense foliage. Adjust spray volume used to plant size.

INSTRUCTIONS

Repeat applications at 4 to 8 day intervals only as needed to control infestations. Applications can be made up to the day of collection.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[At the first sign of insects, in late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind.](#)