

OUTDOOR PET AND DOMESTIC LIVESTOCK USES - PESTS ON SWINE AND POULTRY - BLOWFLIES ETC.

General Information

HOW TO MIX AND APPLY

Unless otherwise directed, mix the recommended amount of HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS per 4 gallons of water and apply in accordance with specific Directions For Use.

Where appropriate, calculate the area to be treated by measuring the length and the width then multiply together to determine the square footage of the area to be treated.

Unless otherwise directed, apply HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS with a pump-up or other commercial type sprayer capable of delivering a coarse spray. When mixing, first fill the sprayer with half of the required amount of water, then measure and add the appropriate amount of HI-YIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS. Complete filling the sprayer with the remaining amount of required water. Shake the sprayer or initiate agitation to mix the product and water. Thorough spray coverage is necessary, but avoid spray runoff.

For large area coverage, this product may also be used through a hose-end type sprayer such as a ferti lome Air-O-Matic Hose-end Spray. Always follow the directions provided with the hose-end sprayer type for proper dilution and application of HIYIELD INDOOR/OUTDOOR BROAD USE INSECTICIDE INCLUDING TURF, GARDENS & ORNAMENTALS.

Do not use food utensils, such as measuring spoons or cups, for food purposes after use in measuring pesticides.

Limitations, Restrictions, and Exceptions

OUTDOOR PET AND DOMESTIC LIVESTOCK USES

FOR USE ONLY ON DOGS AND DOMESTIC LIVESTOCK

ANIMALS AND ANIMAL PREMISE USE CATTLE BARNs, SWINE HOUSES, MILK ROOMS, KENNELS, POULTRY HOUSES, HORSE BARNs, DAIRIES, MILKING PARLORS, STABLES, PADDOCKS.

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 ounces per gallon) (1/2%). For normal infestations dilute (1) part concentrate in 100 parts water (1 1/3 ounces per gallon of water) (8 teaspoonfuls per gallon)(0.1 %).

PESTS ON SWINE AND POULTRY (Do not ship swine for slaughter within 5 days of last treatment)

Spray, dip or sponge to apply 1 pint per pig, especially around ears.

Method

[Dip](#)

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)