

AQUATIC AND OTHER NONCROP SITES - PERENNIAL WEEDS (COGONGRASS)

General Information

GENERAL INFORMATION

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of many herbaceous or woody plants. It may be applied through most standard industrial or field-type sprayers after dilution and through mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most Annual Weeds occur within 2 to 4 days, but on most Perennial Weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on the label, delay application until vegetation has emerged and reached the states described for control of such vegetation under the WEEDS CONTROLLED section of the label. Unemerged plants arising from unattached underground rhizomes or root stocks of Perennials will not be affected by the herbicide and will continue to grow.

For this reason, best control of most Perennial Weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to Annual or Perennial Weeds that have been mowed, grazed, or cut, and have not been allowed to regrow

to the recommended state for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in the labeling. Mixing this product with herbicides or other materials not recommended on the label may result in reduced performance.

For best results, spray coverage should be uniform and complete.

Do not spray weed foliage to the point of runoff.

The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate(s).

Limitations, Restrictions, and Exceptions

WEEDS CONTROLLED

PERENNIAL WEEDS

Apply this product as follows to control or destroy most vigorously growing Perennial Weeds. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

Add 2 or more quarts of non-ionic surfactant per 100 gallons of spray solution to the rates of this product given in this list. Refer to the MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS section for specific uses and application instructions.

Cogongrass: Apply when Cogongrass is at least 18 inches tall and actively growing in late Summer or Fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Method

[Broadcast](#)

Rates

[field rates 0](#)

-

Timings

[Postemergence \(Weed\)](#)