

ALFALFA GROWN FOR SEED - COMMON GROUNDSEL (WASHINGTON)

General Information

GENERAL INSTRUCTIONS AND INFORMATION

APPLICATION INSTRUCTIONS

Do not apply ASULOX Herbicide through any type of irrigation systems.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Limitations, Restrictions, and Exceptions

ALFALFA GROWN FOR SEED

INSTRUCTIONS

Apply the specified rate of Asulox using ground application equipment in a minimum of 15 gallons of water per acre. Thorough and uniform spray coverage on the target weed is very important for best results.

Apply on seedling alfalfa that has at least 3 trifoliolate leaves and any time on established alfalfa including growth from stubble.

Make post-emergence applications of Asulox on actively growing weeds. Apply on small weeds less than 3 inches tall for best results.

Addition of a nonionic surfactant (NIS) at 0.25% by volume (1 quart per 100 gallons) is recommended. Addition of other types of spray adjuvants (such as crop oil concentrates) or fertilizers (such as urea ammonium nitrate) is not recommended.

Asulox causes a slow kill of susceptible weed species and a few weeks following treatment may be required for complete senescence especially under conditions of low temperature and low humidity. Cultural practices that disturb the root system of

targeted weed species may result in less than optimum control with Asulox herbicide. These practices are not recommended within 7 days prior to or within 7 days after applications of Asulox.

RESTRICTIONS

Do not apply more than 7.2 pints of Asulox per acre per growing season.

Do not apply Asulox more than twice per growing season. Allow at least 30 days between applications.

For use under this SLN registration do not apply this product using aerial application equipment or through any type of irrigation system.

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Weed\)](#)

[At least 3 trifoliate leaves.](#)