

ESTABLISHED PERENNIAL GRASSES GROWN FOR SEED - PERENNIAL RYEGRASS, TALL FESCUE (WASHINGTON)

General Information

GENERAL INFORMATION

SOIL TEXTURE: As used on the label, “Coarse soils” are sand, loamy sand or sandy loam soils. “Medium soils” are loam, silt loam, silt, sandy clay, or sandy clay loam. “Fine soils” are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not allow sprays to drift on to adjacent desirable plants.

Apply this product only as specified on the label.

Do not use on other crops grown for food or forage. Observe all cautions and limitations on labeling of all products used in mixtures.

Do not rotate any crop not listed on the label for 18 months following application of DIMETRIC DF 75%.

For all uses: Low-pressure and high-volume hand-wand equipment is prohibited.

GROUND APPLICATION: Apply the proper rate of DIMETRIC DF 75% in a minimum of 10 to 40 gallons of spray mixture per acre broadcast.

Banded Application: Use proportionally less DIMETRIC DF 75% per acre in a band versus a broadcast application. For band application use 1/4 to 1 gallon of spray mix per inch of band width regardless of row spacing.

AERIAL APPLICATION: Where permitted, apply specified rate in a minimum of 2 to 10 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

NOTE: Do not apply aerially when DIMETRIC DF 75% is tank-mixed with Lasso.

For All Applications of DIMETRIC DF 75%: Sprayer must be accurately calibrated before applying METRICDF. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases. Avoid over-application, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops, such as cole crops, onions, peas, or strawberries are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

APPLICATION: Uniform application is essential for satisfactory weed control. Accurate calibration of fertilizer application equipment is essential for uniform distribution to the soil surface. The correct method of application is to apply 1/2 the specified rate and overlap 50 percent or to double apply by splitting the middles to obtain the best distribution pattern. If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

INCORPORATION AND COMBINATION USES: When DIMETRIC DF 75% is to be used in combination with another herbicide, follow directions on the label for combinations, rates, crops, incorporation, and special precautions.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

ESTABLISHED PERENNIAL GRASSES GROWN FOR SEED

For Weed Control in Established Perennial Grasses Grown for Seed in Washington

Used as directed below, Dimetric DF 75% will reduce competition from volunteer seedlings of the indicated crop, annual Bromus species, annual ryegrass, and annual bluegrass. Dimetric DF 75% will control rattail fescue, henbit, ivyleaf speedwell, chickweed, mustards, and shepherdspurse. Control of the volunteer crop and grassy weeds may be enhanced by adding wetting agents containing crop oil.

Follow the directions for use and rates specified on the wetting agent label.

REMARKS

Apply specified dosage as a broadcast spray in at least 15 gallons of spray solution per acre when volunteer grasses are in the 1 to 2 leaf growth stage following full rainfall or irrigation and before active spring growth. Applications made after mid-February may result in excessive crop injury and/or failure to control weeds.

Allow at least 120 days between application and harvest for seed.

Application Restrictions:

1. Apply only to established grasses that are at least one year old and has been harvested for seed at least once.
2. Do not apply to a crop that is under stress, for example disease, severe insect damage, nutrient deficiency, cool to cold temperatures, or deficient or excessive moisture.
3. Do not apply more than once per year.
4. Do not apply this product through any type of irrigation system.
5. Do not apply to bluegrass grasses grown for seed on sand or loamy sand soils. On bluegrass, use 0.2 to 0.25 lbs of Dimetric DF 75% per acre if mixing with another herbicide.
6. Use only on soils with organic matter greater than 1.5% and a soil pH lower than 7.5.

Feeding Restrictions: The crop and crop residues may be fed to livestock or used as bedding.

If the seed crop is terminated and grazed or cut for forage, allow at least 28 days between application and use as animal feed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[When volunteer grasses are in the 1 to 2 leaf growth stage following full rainfall or irrigation and before active spring growth.](#)